

NEON AND [C II] 158 μ m EMISSION LINE PROFILES IN DUSTY STARBURSTS AND ACTIVE GALACTIC NUCLEIANAHIT SAMSONYAN¹, DANIEL WEEDMAN², VIANNEY LEBOUTEILLER³, DONALD BARRY², AND LUSINE SARGSYAN²*Draft version August 9, 2016*

ABSTRACT

The sample of 379 extragalactic sources is presented that have mid-infrared, high resolution spectroscopy with the *Spitzer* Infrared Spectrograph (IRS) and also spectroscopy of the [C II] 158 μ m line with the *Herschel* Photodetector Array Camera and Spectrometer (PACS). The emission line profiles of [Ne II] 12.81 μ m, [Ne III] 15.55 μ m, and [C II] 158 μ m are presented, and intrinsic line widths are determined (full width half maximum of Gaussian profiles after instrumental correction). All line profiles together with overlays comparing positions of PACS and IRS observations are made available in the Cornell Atlas of *Spitzer* IRS Sources (CASSIS). Sources are classified from AGN to starburst based on equivalent widths of the 6.2 μ m polycyclic aromatic hydrocarbon feature. It is found that intrinsic line widths do not change among classification for [C II], with median widths of 207 km s⁻¹ for AGN, 248 km s⁻¹ for composites, and 233 km s⁻¹ for starbursts. The [Ne II] line widths also do not change with classification, but [Ne III] lines are progressively broader from starburst to AGN. A small number of objects with unusually broad lines or unusual redshift differences in any feature are identified.

Subject headings: infrared: galaxies — galaxies: starburst— galaxies: active— galaxies: distances and redshifts—

1. INTRODUCTION

Identifying and understanding the initial formation of massive galaxies and quasars in the early universe is a fundamental goal of observational cosmology. A rapidly developing capability for tracing luminosity sources to high redshifts is the observation of the [C II] 158 μ m emission line at redshifts $z > 4$ using ground based submillimeter interferometers (e.g. Huynh et al. 2013; Wang et al. 2013; Carilli et al. 2013; Banados et al. 2015), with detections now having been made to $z = 7$. This has long been known as the strongest far-infrared line in most sources (Stacey et al. 1991; Malhotra et al. 1997; Nikola et al. 1998; Luhman et al. 2003; Brauher et al. 2008), often carrying $\sim 1\%$ of the total source luminosity, and is thought to be associated with star formation because it should arise within the photodissociation region (PDR) surrounding starbursts (Tielens and Hollenbach 1985; Helou et al. 2001; Malhotra et al. 2001; Meijerink et al. 2007).

In many high redshift sources, especially dust obscured sources, the [C II] feature is the only diagnostic atomic emission line which can be observed. This makes it vital to optimize use of this line for learning about intrinsic source properties. Not only line luminosities but also accurate line profiles are observed. For example, the submillimeter galaxy in the group BRI 1202-0725 at $z = 4.7$ has a [C II] FWHM of 700 km s⁻¹ measured with the Atacama Large Millimeter Array, but the associated quasar has FWHM of only 275 km s⁻¹ (Carilli et al. 2013). What can we learn from such differences?

A great amount of diagnostic information concern-

ing the observable differences between dusty, obscured sources powered by active galactic nuclei (AGN) and those powered by rapid star formation (“starbursts”) accumulated with the mid-infrared spectroscopy of the Infrared Spectrograph (IRS; Houck et al. 2004) on the *Spitzer* Space Telescope (Werner et al. 2004). Classifications of AGN and starburst sources were developed using both the strength of PAH features in low resolution spectra and various emission line ratios in high resolution spectra (e.g. Brandl et al. 2006; Farrah et al. 2007; Desai et al. 2007; Bernard-Salas et al. 2009; Weedman and Houck 2009; Veilleux et al. 2009; Tommasin et al. 2010; Wu et al. 2010; Sargsyan et al. 2011; Stierwalt et al. 2013; Inami et al. 2013).

More recently, numerous observations of the [C II] line were made (Sargsyan et al. 2012; Farrah et al. 2013; Diaz-Santos et al. 2014; Sargsyan et al. 2014; De Looze et al. 2014; Diaz-Santos et al. 2013) using the Photodetector Array Camera and Spectrometer (PACS) instrument (Poglitsch et al. 2010) on the *Herschel* Space Observatory (Pilbratt et al. 2010) and included many sources that had previously been observed by the IRS. Our own previous studies were specifically designed to compare the [C II] line luminosities to mid-infrared polycyclic aromatic hydrocarbon (PAH) emission features and mid-infrared emission lines observed with the IRS in order to calibrate the star formation rate (SFR) using the [C II] line. We found that [C II] luminosities most closely track [Ne II] luminosities, from which we concluded that the [C II] is showing PDRs associated with the HII regions seen in [Ne II].

The observational study of forbidden emission line profiles has long been a key technique for classifying and understanding active galaxies. The most extensive early studies used the optical [O III] line at 0.50 μ m, from which a conclusion was reached that the width of this line could generally distinguish AGN and starbursts

¹ Byurakan Astrophysical Observatory, Byurakan, Armenia, anahit.sam@gmail.com

² Cornell Center for Astrophysics and Planetary Science, Cornell University, Ithaca, NY 14853; dweedman@astro.cornell.edu

³ Laboratoire AIM, CEA/DSM-CNRS-Universite Paris Diderot, DAPNIA/Service d’Astrophysique, Saclay, France

(Feldman et al. 1982; Whittle 1985). All conclusions based on optical lines were restricted to sources with little extinction by dust, however, which led to uncertainties in interpreting differences among profiles if different components of sources have different values of extinction. Now that forbidden emission line profiles are available in mid-infrared and far infrared, comparisons can be made without this uncertainty. An initial indication that such line profiles provide interesting results is in the line profile comparisons by Spoon et al. (2013) among 24 ultraluminous infrared galaxies (ULIRGS) with the goal of seeking outflows from AGN.

To evaluate further the utility of the [C II] line for understanding AGN and starbursts, we compare in the present paper the [C II] emission line profiles to the mid-infrared diagnostics for all extragalactic sources which have archival observations available for both PACS [C II] and IRS spectra. This study of line profiles is an extension of previous analyses of total [C II] line fluxes. We present emission line profiles for [C II], [Ne II] 12.81 μm , and [Ne III] 15.55 μm for extragalactic sources which have archival observations available for both PACS [C II] and IRS high resolution spectra. Our resulting sample of 379 extragalactic sources includes all sources with both [C II] and IRS high resolution observations, except for nearby extended galaxies for which IRS observations were made in a mapping mode. The latter are omitted because our goal is to understand overall, global properties of sources that would generally be unresolved at high redshift. There are 390 [C II] profiles in our total sample, representing 379 different IRS pointings (the remaining 11 are duplicate [C II] observations made at the same IRS position).

We present the source list and measurements of line widths and redshifts along with overall comparisons among line widths. All of the [C II] line profiles with Gaussian fits are made available in the Cornell Atlas of *Spitzer*/IRS Sources (CASSIS; Lebouteiller et al. 2011, 2015⁴). Further comparisons with other source properties will be considered in a future paper (in preparation, Samsonyan 2016).

2. OBSERVATIONS

2.1. Sample Selection

Our sample was selected by examining abstracts of *Herschel* observing programs which described observations of extragalactic emission line sources⁵ and then studying archival sources from these programs to identify those for which the [C II] line was observed, either in line or range spectroscopy. Once all extragalactic sources with [C II] observations were identified, we searched for these sources in CASSIS to locate sources having high resolution IRS observations, necessary for line profile information. The individual *Herschel* observing programs and the number of sources taken from each are SDP-esturm-3, DDT-esturm-4, and KPGT-esturm-1(56); KPGT-smadde01-(13); KPOT-pvanderw-1(11); GT1-lspinogl-4 and GT2-lspinogl-6(14); OT1-dfarrah-1 and OT2-dfarrah-5(23); OT1-dweedman-1(112); OT1-

larmus-1(136); OT2-tdiazsan-1(4); OT1-lyoung-1(5); OT1-nwerner-1(3); OT1-sveilleu-2(1); OT2-idelooze-1(1); OT2-lsargsyan-1(2); OT2-nnesvad-3(2); OT2-pguillar-7(5); KPOT-aedge-1(1); and OT1-bweiner-1(1). Although we intended a complete search, it is possible that some sources having both [C II] and IRS high resolution observations have been overlooked. The source list and all results are given in Table 1.

2.2. IRS High Resolution Spectra

A few previous studies have determined line widths from IRS high resolution spectra (Dasyra et al. 2011; Spoon et al. 2013; Inami et al. 2013). These results have all been approximated by applying a uniform estimate of instrumental resolution using the typical resolution of $\sim 500 \text{ km s}^{-1}$ for full width half maximum (FWHM) of the instrumental profile listed in the instrument description. For our analysis, we apply two additional steps for improving determination of the instrumental profile. The first is to determine empirical instrumental profiles for individual emission lines, and the second is to apply an “optimal” extraction for the high resolution spectra, which enhances signal to noise (S/N) for unresolved sources. These steps are described below.

Spectral extraction is the process of producing a one dimensional spectrum from the two dimensional images obtained with the IRS detectors. The best possible extraction of unresolved sources applies the point spread function (PSF) of the telescope/spectrograph combination so that the pixels in the cross dispersion direction are weighted by the fraction of source flux which falls on them, thereby reducing background and instrumental noise. This is defined as an “optimal” extraction. These extractions have been implemented for the spectra shown in CASSIS using empirically determined PSFs for both low resolution (Lebouteiller et al. 2011) and high resolution (Lebouteiller et al. 2015). If a source is fully extended over the observing aperture, a better alternative is a full aperture extraction that equally weights all pixels within the aperture. The choices provided in CASSIS for spectral extractions of high resolution IRS spectra are described in detail by Lebouteiller et al. (2015). All previous measures of IRS high resolution spectra have used full aperture extractions, which are provided by the “post-BCD” spectra of the *Spitzer* Heritage Archive⁶.

For both low resolution and high resolution spectra, the source extent is estimated in CASSIS by noting differences between the source spatial profile compared to the PSF. This estimate is more reliable for low resolution spectra than for high resolution because of the limited spatial profile sampling in high resolution; the PSF has full width half maximum (FWHM) of about 2 pixels compared to the 6 pixel length of the high resolution slits. The majority of our sources are not extended on the scale of the observing aperture, so our final adopted measurements utilize the optimal extraction. For our objective of measuring line profiles, the best possible S/N is crucial in order to obtain the best profiles. This also leads to the choice of optimal extraction even if some flux is underestimated for extended sources.

The extraction labeled as “optimal” in CASSIS uses a simultaneous fitting to the two separate observing nod

⁴ <http://cassis.sirtf.com>. The [C II] profiles are in <http://cassis.sirtf.com/herschel>. CASSIS is a product of the Infrared Science Center at Cornell University.

⁵ <http://www.cosmos.esa.int/web/herschel/observations>

⁶ <http://irsa.ipac.caltech.edu/data/SPITZER/docs/spitzerdataarchives>

positions. This option also provides an estimate of the underlying background by using residuals beneath the fitted PSF. Another alternative, “optimal differential”, that differences the two nods before fitting the PSF profile is also given in CASSIS for comparison to aid in removing bad pixels, but this method gives a reliable flux calibration only for unresolved sources. CASSIS illustrates both of the optimal extractions in addition to a full aperture extraction so that users can make a final choice but recommends “optimal” if the source appears spatially unresolved.

The IRS instrumental spectral resolution for the high resolution modes is approximately defined by the two pixel projected width of the observing aperture. Spectral resolution varies among different lines because the IRS high resolution spectrographs are echelle spectrographs with different lines seen in different orders, so resolution increases as the order increases. Because our objective is the best possible measurement of line widths, the instrumental resolutions for the [Ne II] and [Ne III] lines presented in this paper are measured empirically. This is done using spatially unresolved planetary nebulae in which these emission lines are strong. The intrinsic widths of planetary nebula emission lines are much smaller than the IRS instrumental line width so the observed line profiles illustrate the instrumental profile.

The sources used are the planetary nebulae from the Large Magellanic Cloud, Small Magellanic Cloud, and Galactic Halo included in Bernard-Salas et al. (2009b) and Pottasch and Bernard-Salas (2010). Because these planetaries are spatially unresolved, the CASSIS optimal extraction is used to measure the lines. Gaussian fits match the observed profiles well so we adopt instrumental profiles that are Gaussians described by the FWHM of the planetary nebula profiles.

The FWHM and dispersion among individual planetaries is $\text{FWHM}([\text{Ne II}]) = 331 \pm 35 \text{ km s}^{-1}$ and $\text{FWHM}([\text{Ne III}]) = 377 \pm 35 \text{ km s}^{-1}$. The somewhat better resolution for [Ne II] $12.81 \mu\text{m}$ is expected, because this line is in short-high order 16 whereas [Ne III] $15.55 \mu\text{m}$ is in order 13. These instrumental line widths were also measured using full aperture extractions; the full aperture extractions give consistent but slightly larger median widths compared to the optimal extractions, 371 km s^{-1} for [Ne II] and 465 km s^{-1} for [Ne III]. These differences can be explained by small tilts of the observing slits compared to the dispersion direction of the spectra. The optimal extractions give more weight to the center of the slit where any line smearing from tilt effects is negligible, but the full extractions equally weight flux near the edges, where slit tilts would produce a larger shift of the dispersed image.

Because the optimal extraction is used for sources in this paper, we adopt as the instrumental resolution profile (FWHM_{res}) the values of 331 km s^{-1} for [Ne II] and 377 km s^{-1} for [Ne III], with an uncertainty of about 10% for each. The intrinsic $\text{FWHM}_{\text{intrinsic}}$ of the observed spectral lines in sources are then determined as $\text{FWHM}_{\text{intrinsic}}^2 = \text{FWHM}_{\text{observed}}^2 - \text{FWHM}_{\text{res}}^2$.

Redshift measurement for sources depend on the position of the line centroids, which depends on the wavelength calibration of the spectra (“WAVESAMP” files). This calibration is applied differently in optimal and full

aperture extraction. To determine any systematic uncertainties arising from the extraction technique, we compared line centroids for the planetary nebulae and find that the 1σ dispersion between full aperture and optimal extractions for the Neon lines is $\pm 25 \text{ km s}^{-1}$. These results mean that redshift uncertainties arising from different analysis techniques for the high resolution spectra are only ± 0.00008 .

The observed results for line widths from the CASSIS optimal extractions are given in Table 1. All IRS emission lines, including those obtained with the alternative extractions, can be examined within CASSIS by source name or coordinate using the browse tools provided without requiring any downloads of the spectra. The CASSIS results provide the actually observed coordinate of the source, which sometimes differs from the nominal coordinate corresponding to the source name in other data bases. In “cluster” mode for IRS observations, several independent sources may be included within the same Astronomical Observation Request (AOR), and these sources are sometimes components of the same galaxy. For these reasons, it is necessary to use the accurate coordinates from CASSIS for comparisons to locate the PACS spaxel corresponding to the same position, and these coordinates are those in Table 1.

To determine uncertainties in FWHM for line profiles depending on extraction technique, we compared both full aperture and optimal extractions for the [Ne II] and [Ne III] lines in all sources. Line profiles were measured using Gaussian fits to the [Ne II] and [Ne III] emission lines with a first order underlying continuum, fitting with the Spectroscopic Modeling Analysis and Reduction Tool (SMART; Higdon et al. 2004, Lebouteiller et al. 2010). The dispersions among the differences between full aperture and optimal extractions for the same profiles are a measure of the observational uncertainties that arise from independent fitting of undersampled profiles with different Gaussians. For both lines, these dispersions ($\pm 50 \text{ km s}^{-1}$ for [Ne II] and $\pm 46 \text{ km s}^{-1}$ for [Ne III]) are $\sim 10\%$ of the observed median line width (470 km s^{-1} for [Ne II] and 512 km s^{-1} for [Ne III]).

Dispersions are a larger fraction of the measured width for narrower lines so the fractional uncertainty is greater for narrow lines. This effect is an additional reason to prefer optimal extractions for the line measurements. In some cases, the observed FWHM from the best Gaussian fit is smaller than the instrumental resolution, which cannot be physically correct. In these cases, the adopted intrinsic FWHM is arbitrarily listed as 100 km s^{-1} and displayed at that value in Figures so that these anomalous cases can be recognized. There is an offset in measured FWHM such that the full aperture measures are systematically larger, by 20 km s^{-1} for [Ne II] and 70 km s^{-1} for [Ne III]. We attribute this difference to a similar effect as noted above for instrumental resolution - that weighting all pixels of the tilted slit evenly in the full aperture extraction leads to a smearing of the line in direction of dispersion, which artificially increases the line width.

2.3. *Herschel* PACS [C II] Spectra

The PACS instrument (Poglitsch et al. 2010) simultaneously obtains spectra at 25 positions in a source with square apertures called “spaxels”. If all sources were




FIG. 1.— Example of PACS spaxels compared to IRS high resolution slits, for source Markarian 18. Squares are spaxels of size $9.4'' \times 9.4''$; large rectangle is IRS long-high slit, and small rectangle is IRS short-high slit. Crosses show coordinate of source in IRS slits. All Neon lines measured are in the short-high slit. The closest spaxel to position of the IRS slit is identified in Table 1; numbering convention used in Table 1 for rows and columns of spaxels is that upper left is 5,5; lower left is 5,1; upper right is 1,5; and lower right is 1,1. E is to the left and N to the top of the image. These overlays for all sources are shown in CASSIS at <http://cassis.sirtf.com/herschel>.

spatially unresolved and all pointings were perfect, direct comparisons of lines observed with IRS and PACS would be straightforward. Although neither condition is perfectly met for all of these observations, both the IRS slit and a PACS spaxel observe similar fractions of a source. The IRS short wavelength, high resolution observing slit for the Neon lines is $4.7''$ wide, designed to match approximately the FWHM of the telescope diffraction profile at these wavelengths. The PACS spaxels are $9.4''$ wide, also nearly equal to the FWHM of the *Herschel* diffraction profile at the [C II] wavelength. Because of this similarity in fraction of a source observed by the two spectrographs, we compare line profiles seen in the IRS slit to the [C II] profile seen in the single PACS spaxel that is closest in location to the IRS slit, as illustrated in Figure 1.

A factor which might affect profile widths in spatially resolved sources is the size of the emitting region observed by PACS. These sizes vary by a large amount because distances range from 0.05 Mpc to 1003 Mpc. In Table 1, we list the projected sizes of the PACS $9.4''$ spaxel based on the angular size distances to the sources. Redshift independent distances are used if these are listed in the NASA/IPAC Extragalactic Database (NED). If not, redshift derived distances are determined from Wright (2006) using $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.27$ and $\Omega_\Lambda = 0.73$. The distribution in projected spaxel size among sources in Table 1 is shown in Figure 2. The distributions illustrate that the PACS results refer primarily to global properties of sources on observed scales of many kpc.

For our studies of line profiles, an important consideration is to match the position of a PACS spaxel to the position of the IRS aperture. We did this individually for each source, producing overlays of PACS spaxels and IRS aperture using the observed positions given in




FIG. 2.— Distributions in projected size of $9.4''$ PACS spaxel among all sources in Table 1. Histograms show actual number of sources of different classifications in each bin of width one kpc.




FIG. 3.— Example of a [C II] profile produced by PACSman and displayed in CASSIS, for source Markarian 18. The fit by PACSman produces the FWHM and uncertainty of the observed Gaussian profile (solid line) fit to the data points (broken line and error bars), together with line flux and equivalent width. The line profile is shown at the [C II] redshift given in Table 1, corrected to the local standard of rest. Any assumed instrumental width (FWHM-th; dashed line) may be input to yield intrinsic width (Broadening) after removing this instrumental width from the observed profile. Instrumental FWHM used for all sources in this paper is 236 km s^{-1} , determined empirically from observations of 30 Doradus, so intrinsic widths given in Table 1 are determined as $\text{FWHM}_{\text{intrinsic}}^2 = \text{FWHM}_{\text{observed}}^2 - 236^2$. All profiles used in this paper and the observed FWHM are illustrated in CASSIS at <http://cassis.sirtf.com/herschel>, although slightly different values of FWHM-th are sometimes shown for illustration. For noisy profiles with an observed FWHM fit of $< 236 \text{ km s}^{-1}$, an artificial value of FWHM-th is shown in the profiles so that the value determined for “broadening” remains positive. In Table 1, all profiles with observed FWHM $< 236 \text{ km s}^{-1}$ are arbitrarily assigned values of 100 km s^{-1} for intrinsic width.

the headers for the final IRS spectra, the coordinates in Table 1. Overlays such as the example in Figure 1 are shown in CASSIS for all of our sources. The individual spaxel corresponding most closely to the position of the IRS aperture is listed in Table 1, using the numbering convention shown in Figure 1, and the profile from this spaxel is the [C II] profile shown in CASSIS.

All [C II] observations were reduced with version 12.1.0 of the *Herschel* Interactive Processing Environ-

ment (HIPE), together with the “PACSman” software (Lebouteiller et al. 2012) to fit line profiles and continuum within the spaxels. An illustration of a Gaussian line fit provided by PACSman is in Figure 3, and all profiles which we use are shown for the individual sources in CASSIS, searchable by source name or coordinate. Displaying Gaussian fits is particularly useful for seeking sources which might have a core Gaussian but broad or asymmetric underlying wings (e.g. Janssen et al. 2016). There are 34 sources in our sample for which [C II] was observed but yields a profile with insufficient S/N for an accurate width or radial velocity measurement. These profiles are shown as “no data” in Table 1, but they are illustrated in CASSIS with an attempted Gaussian profile fit at the line position.

As was done for the IRS spectra, we determined an empirical instrumental profile for the [C II] observations. This was done using all 25 spaxels in the observation of the 30 Doradus HII region listed in Table 1, because all spaxels give the same observed FWHM, the intrinsic line widths are small compared to the instrumental resolution, and the intrinsic line widths are known. The observed FWHM among these 25 profiles is 239 km s^{-1} . Internal motions within the HII region have been measured for the optical [O III] $0.50 \mu\text{m}$ line and have median FWHM of 45 km s^{-1} FWHM at spatial resolutions similar to PACS (Smith and Weedman 1972). Removing this intrinsic width from the width observed by PACS results in an instrumental FWHM of 236 km s^{-1} . The intrinsic FWHM_{intrinsic} of the observed [C II] lines in sources are then determined as $\text{FWHM}_{\text{intrinsic}}^2 = \text{FWHM}_{\text{observed}}^2 - 236^2$. The intrinsic widths are listed in Table 1.

3. ANALYSIS OF [C II] AND NEON LINE PROFILES

Our primary objective for the present paper is to present the measurements in Table 1 and the profiles in CASSIS, which together provide an observational atlas of [C II] $158 \mu\text{m}$, [Ne II] $12.81 \mu\text{m}$, and [Ne III] $15.55 \mu\text{m}$ line widths that can be used for comparing among various properties for a wide variety of sources. Our eventual objective is to understand in more detail what determines line widths in different galaxies. For now, we only use these results to determine if there are trends with classification as AGN or starburst, or any systematic differences among [C II] and mid-infrared Neon lines.

3.1. Comparisons of Line Profile Widths

Previous studies using IRS line widths found some indications that higher ionization lines sometimes show broader widths or outflow effects from AGN, but these effects were found primarily for the high ionization [SIV] and [OIV] lines in a few cases of sources with the broadest lines (Dasyra et al. 2011; Spoon et al. 2013; Inami et al. 2013). We emphasize the lower ionization Neon lines, especially [Ne II] because this line most closely tracks [C II] in comparisons of line fluxes (Sargsyan et al. 2014). These Neon lines are also the best for comparison with previous studies of line profiles of the strongest optical forbidden line, [O III] $0.50 \mu\text{m}$ (Feldman et al. 1982; Whittle 1985; Nelson and Whittle 1996), because of similar ionization potentials (21.6 eV to produce Ne II, 41.0 eV to produce Ne III, and 35.1 eV to produce O III.)

The AGN/starburst classification used here, based on strength of the $6.2 \mu\text{m}$ PAH feature, is similar to

that used in many previous studies, although different authors use different PAH features and different methods for measuring strength (e.g. Genzel et al. 1998; Laurent et al. 2000; Brandl et al. 2006; Desai et al. 2007; Veilleux et al. 2009; Tommasin et al. 2010; Wu et al. 2010; Sargsyan et al. 2011; Petric et al. 2011; Stierwalt et al. 2013). The essential result used by all of these authors is that the PAH features increase in strength as the starburst component increases although different authors adopt somewhat different definitions of AGN-dominated compared to starburst-dominated. The quantitative divisions illustrated in our results are that AGN have $\text{EW}(6.2 \mu\text{m}) < 0.1 \mu\text{m}$, composite AGN plus starburst have $0.1 \mu\text{m} < \text{EW}(6.2 \mu\text{m}) < 0.4 \mu\text{m}$, and starbursts have $\text{EW}(6.2 \mu\text{m}) > 0.4 \mu\text{m}$.

The divisions we use began with an IRS spectroscopic survey of starburst galaxies by Brandl et al. (2006) and in a flux limited IRS survey of sources including all classifications (Houck et al. 2007), subsequently verified by comparison with optical classifications using several hundred IRS spectra (Sargsyan et al. 2011). Our primary motive for using the $6.2 \mu\text{m}$ feature instead of the stronger $11.3 \mu\text{m}$ feature has been to allow classification of sources having sufficiently high redshifts that the $11.3 \mu\text{m}$ feature is not visible in IRS spectra. The strongest PAH complex centered at $7.7 \mu\text{m}$ is not used because this feature can be confused with an apparent spectral peak at similar wavelength arising because of strong absorption on either side of the peak in heavily obscured AGN. By illustrating various parameters regarding line profiles as functions of $\text{EW}(6.2 \mu\text{m})$, any trends with AGN/starburst classes are readily seen.

The well established correlation between level of ionization and PAH classification of AGN and starbursts is illustrated in Sargsyan et al. (2014) for various flux ratios among mid-infrared and [C II] emission lines as a function of PAH classification. For our present use, the important result is that the [Ne II] line tracks [C II] independent of classification whereas the higher ionization [Ne III] line is stronger relative to [C II] in AGN than in starbursts. (Ratio of line fluxes for [Ne III]/[Ne II] for the optimal extractions is given in Table 1 for each source.) This implies that the line profiles from [Ne III] should be more affected by the AGN, whereas the [Ne II] and [C II] lines should characterize the starburst component. We first consider, therefore, whether line profiles for these three features show any differences between AGN and starbursts.

Comparisons of AGN/starburst classification with intrinsic line widths (those given in Table 1, after correction for instrumental broadening) are shown in Figures 4, 5, and 6. We think that the most important result is shown in Figure 4 for [C II] because a primary goal of our study is to seek any evidence that the [C II] line width is a diagnostic of AGN/starburst classification. There is no evidence of any trend, however. The medians and dispersions among [C II] line widths in Figure 4 are very similar for all classifications, with medians 207 km s^{-1} for AGN, 248 km s^{-1} for composites, and 233 km s^{-1} for starbursts with dispersions in all cases of about $\pm 130 \text{ km s}^{-1}$. The most important result for [C II] in Figure 4, therefore, is that line widths do not change among classification, indicating that [C II] is dominated by the starbursts within

any source and not affected by the presence of an AGN. This confirms our conclusions in Sargsyan et al. (2012) and Sargsyan et al. (2014) based only on line fluxes.

The results for [Ne II] in Figure 5 show a similar result to [C II], with only a slight trend of increased widths for AGN; the median FWHM for AGN and composites is 340 km s^{-1} compared to 300 km s^{-1} for starbursts. The best comparison with [C II] is for the starbursts to rule out any broadening of [Ne II] by an AGN. For starbursts, the difference between median FWHM of 233 km s^{-1} for [C II] and 300 km s^{-1} for [Ne II] is interesting, but we cannot be confident that it is meaningful. This difference is less than the uncertainty in the individual IRS width measures shown in Figure 5. The difference would disappear if we assume a broader instrumental line width for [Ne II] than we have adopted, requiring an instrumental width of 380 km s^{-1} instead of the 331 km s^{-1} that is used. This larger value is close to the full aperture instrumental width discussed in section 2.2, but such a large instrumental width would not be appropriate for unresolved sources, which require an optimal extraction. If this larger instrumental width were used, full aperture extractions would also have to be used, and these are systematically larger than the optimal extraction widths that are given in Table 1. The net result would remain that intrinsic widths for [Ne II] are greater than those for [C II]. We must conclude, therefore, that we cannot precisely compare the [C II] widths from PACS with the Neon widths from the IRS because of uncertainties in determining the most accurate extraction procedure for the IRS lines.

Despite these uncertainties among comparisons of [C II] and Neon line widths, the relative widths between the [Ne II] and [Ne III] lines are meaningful. Unlike the [Ne II] FWHM in Figure 5, the [Ne III] FWHM in Figure 6 show a trend for increasing line widths from starbursts through AGN. Both the median values of line widths and upper values of the dispersions are progressively larger from starbursts through composites to AGN, with medians of 289 km s^{-1} for starbursts, 367 km s^{-1} for composites, and 426 km s^{-1} for AGN. Upper values of the observed one σ dispersions are 417 km s^{-1} for starbursts, 580 km s^{-1} for composites, and 748 km s^{-1} for AGN. This trend for AGN to be offset to larger values of intrinsic [Ne III] line width is independent of the adopted value for [Ne III] instrumental width and demonstrates that some additional [Ne III] line broadening is associated with AGN. This is additional confirmation that we should not expect similar profiles between [C II] and higher ionization features such as [Ne III] because they often arise in different physical regions of sources.

Having confirmed that [C II] and [Ne II] line profiles generally track one another, an additional use of the results in Table 1 is to compare line widths for [C II] and [Ne II] lines to find anomalous sources that are broad in one line but not the other. For example, are there sources where [C II] arises in regions where [Ne II] is not seen, either because of too much extinction or because there is a diffuse [C II] not associated with starbursts? Or, are there sources in which even the low ionization [Ne II] is dominated by the AGN? This comparison is shown in Figure 7. This Figure is used to identify sources with unusually broad features, defined as having FWHM that




FIG. 4.— Intrinsic FWHM of [C II] $158 \mu\text{m}$ in km s^{-1} compared to $\text{EW}(6.2 \mu\text{m})$ from Table 1. Large crosses show median line width and one σ dispersion of line widths within classifications. Thin vertical lines divide classifications; AGN have $\text{EW}(6.2 \mu\text{m}) < 0.1 \mu\text{m}$, composite AGN plus starburst have $0.1 \mu\text{m} < \text{EW}(6.2 \mu\text{m}) < 0.4 \mu\text{m}$, and starbursts have $\text{EW}(6.2 \mu\text{m}) > 0.4 \mu\text{m}$. Small symbols are sources with measured line widths but upper limits in $\text{EW}(6.2 \mu\text{m})$. Small crosses are sources in which the observed FWHM appears smaller than the instrumental resolution so FWHM arbitrarily assumed as 100 km s^{-1} . Observational uncertainty for [C II] FWHM is typically smaller than size of symbols; uncertainties for each source are displayed in CASSIS.




FIG. 5.— Intrinsic FWHM of [Ne II] $12.81 \mu\text{m}$ in km s^{-1} compared to $\text{EW}(6.2 \mu\text{m})$ from Table 1. Symbols and explanations are as in Figure 4, with additional error bar that shows one σ dispersion of differences between full aperture and optimal extractions of the same line profiles which is the measure of uncertainty adopted for observed FWHM, discussed in section 2.2.

exceed the median values by two σ . These sources are noted in Table 1 for further future study.

A similar comparison is also made in Figure 8 between [C II] and [Ne III], although it would be less surprising in this case to find broad [Ne III] associated with an AGN. The few [C II] sources that equal or exceed the [Ne III] width are particularly interesting. Broad profiles seen in this comparison are also noted in Table 1.

3.2. Comparisons of Radial Velocities

An alternative way to seek sources with significant differences between the physical locations of [C II] and Neon




FIG. 6.— Intrinsic FWHM of [Ne III] 15.55 μm in km s^{-1} compared to $\text{EW}(6.2 \mu\text{m})$ from Table 1. Symbols and explanations are as in Figures 4 and 5.




FIG. 8.— Intrinsic [C II] width compared to [Ne III] width in km s^{-1} from Table 1. Symbols and explanations are as in Figure 7, with error bar showing observational uncertainty of [Ne III] FWHM from Figure 6.




FIG. 7.— Intrinsic [C II] width compared to [Ne II] width in km s^{-1} from Table 1. Small crosses are sources in which the observed FWHM appears smaller than the instrumental resolution so FWHM arbitrarily assumed as 100 km s^{-1} . Thick cross shows median (228 km s^{-1}) and one σ dispersion for [C II], and thin cross shows median (333 km s^{-1}) and one σ dispersion for [Ne II] using sources of all AGN/starburst classifications. Sources outside of box have FWHM that exceed medians by two σ . Observational uncertainty for [C II] FWHM is typically smaller than size of symbols; uncertainties for each source are displayed in CASSIS. Error bar is observational uncertainty of [Ne II] FWHM, as in Figure 5.

emission is to compare observed radial velocities measured in these different features. These comparisons are in Figures 9 and 10. The velocity differences dV between the profile centroids are given in Table 1, determined from the redshifts measured independently for the [C II] and Neon lines and transformed to velocity units so that $dV = c[z(\text{[C II]}) - z(\text{Neon})]$. Comparisons of dV are made to the [C II] FWHM to learn if broader lines have systematically more scatter in dV measures, although we see no indication of this. From these plots, sources with dV differing from the medians by more than 2σ are selected and noted in Table 1, because these sources may also prove to be unusual. Dispersions in dV measures between [C II] and Neon are $\sim \pm 50 \text{ km s}^{-1}$, or $\sim 10\%$ of the observed median FWHM for the Neon lines.




FIG. 9.— Difference dV between [C II] radial velocity and [Ne II] radial velocity in km s^{-1} compared to [C II] line width. Large cross shows median (horizontal line) and 1σ dispersion about median (vertical line) of radial velocity differences. Small crosses are sources in which the observed [C II] FWHM appears smaller than the instrumental resolution so FWHM arbitrarily assumed as 100 km s^{-1} . Sources outside of lines have dV that exceed median by two σ .

In both Figures 9 and 10, there is little difference in median radial velocities determined from PACS [C II] or IRS Neon. For $dV([\text{C II}] - [\text{Ne II}])$, the median difference is 21 km s^{-1} and is 34 km s^{-1} for $dV([\text{C II}] - [\text{Ne III}])$. These small systematic differences are a confirmation of the careful wavelength calibrations that were done independently for both instruments. There is a known instrumental effect in PACS which can lead to a skewed profile and offset in velocity. This effect arises when an unresolved source is not perfectly centered in a spaxel⁷ and can produce velocity offsets up to $\sim \pm 30 \text{ km s}^{-1}$ for [C II]. This is less than the offsets which we note as unusually large, and offsets caused by this instrumental effect would be recognized by having a skewed profile.

⁷ <http://herschel.esac.esa.int/Docs/PACS/pdf/pacs-om.pdf>, Figures 4.16 and 4.18




FIG. 10.— Difference dV in [C II] radial velocity compared to [Ne III] radial velocity in km s^{-1} . Explanation of symbols same as in Figure 9.

4. SUMMARY AND CONCLUSIONS

We measure emission line profiles and redshifts for extragalactic sources observed in both [C II] $158 \mu\text{m}$ with *Herschel* PACS together with [Ne II] $12.81 \mu\text{m}$ and [Ne III] $15.55 \mu\text{m}$ observed with the high resolution *Spitzer* IRS. Data are presented and compared for 379 different sources. Results for [C II] and Neon are compared by locating the PACS observing spaxel that most closely corresponds to the position of the IRS slit. Intrinsic line profile widths are determined by applying empirically measured instrumental widths from observed planetary nebulae or HII regions. All [C II] and Neon line profiles together with overlays of PACS spaxels compared to IRS slits are illustrated in the CASSIS spectral atlas (<http://cassis.sirtf.com/herschel>).

Sources are classified as AGN, composite, or starburst based on equivalent width of the PAH $6.2 \mu\text{m}$ feature. The median intrinsic FWHM for [C II] shows no change with classification, being 207 km s^{-1} for AGN,

248 km s^{-1} for composites, and 233 km s^{-1} for starbursts with dispersions in intrinsic line widths of about $\pm 130 \text{ km s}^{-1}$. Results show that [C II] line widths generally match those of [Ne II], as previously indicated in comparisons of line fluxes. A small number of sources are identified with unusually broad lines or with radial velocity differences between [C II] and Neon measures. Accurate redshifts are determined for sources as demonstrated by a systematic difference of only 21 km s^{-1} between the independent measures of [C II] and [Ne II] radial velocities using PACS and IRS.

This paper and the CASSIS atlas of IRS spectra are dedicated to our friend and colleague James R. Houck, who died September 18, 2015. Jim Houck worked for 20 years to develop the Infrared Spectrograph for the *Spitzer* Space Telescope, which has enabled this research and hundreds of other papers based on mid-infrared spectroscopy. Thank you, Jim.

We acknowledge use of archival data from the *Herschel* PACS instrument, which was developed by a consortium of institutes led by MPE (Germany) and including UVIE (Austria); KU Leuven, CSL, IMEC (Belgium); CEA, LAM (France); MPIA (Germany), INAF-IFSI/OAA/OAP/OAT, LENS, SISSA (Italy); and IAC (Spain). This development was supported by the funding agencies BMVIT (Austria), ESA-PRODEX (Belgium), CEA/CNES (France), DLR (Germany), ASI/INAF (Italy), and CICYT/MCYT (Spain). This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. Partial support for this work at Cornell University was provided by NASA through Contracts issued by the NASA *Herschel* Science Center. We thank Dieter Engels and the Hamburger Sternwarte for hospitality during the initial phase of this effort.

REFERENCES

- Banados, E., Decarli, R., Walter, F., Venemans, B., Farina, E., and Fan, X. 2015, *ApJ*, 805, 8
- Bernard-Salas, J., Spoon, H. W. W., Charmandaris, V. et al. 2009, *ApJ*, 672, 274
- Bernard-Salas, J., Peeters, E., Sloan, G. C. et al. 2009, *ApJ*, 699, 1541
- Brandl, B. R., Bernard-Salas, J., Spoon, H. W. W., et al. 2006, *ApJ*, 653, 1129.
- Brauher, J. R., Dale, D. A., and Helou, G. 2008, *ApJS*, 178, 280
- Carilli, C. L., Riechers, D., Walter, F., et al. 2013, *ApJ*, 763, 120
- Dasyra, K. M., Ho, L. C., Netzer, H. et al. 2011, *ApJ*, 740, 94
- De Looze, I., Cormier, D., Lebouteiller, V. et al. 2014, *A&A*, 568, 62
- Desai, V., Armus, L., Spoon, H. W. W., et al. 2007, *ApJ*, 669, 810
- Diaz-Santos, T., Armus, L., Charmandaris, V., et al. 2013, *ApJ*, 774, 68
- Diaz-Santos, T., Armus, L., Charmandaris, V., et al. 2014, *ApJ*, 788, 17
- Farrah, D., Bernard-Salas, J., Spoon, H. W. W., et al. 2007, *ApJ*, 667, 149
- Farrah, D., Lebouteiller, V., Spoon, H. W. W., et al. 2013, *ApJ*, 776, 38
- Feldman, F. R., Weedman, D. W., Balzano, V. A. and Ramsey, L. W. 1982, *ApJ*, 256, 427
- Genzel, R., Lutz, D., Sturm, E., et al. 1998, *ApJ*, 498, 579
- Higdon, S. J. U., Devost, D., Higdon, J. L., et al. 2004, *PASP*, 116, 975
- Helou, G., Malhotra, S., Hollenbach, D. J., Dale, D. A., and Contursi, A. 2001, *ApJ*, 548, L73
- Houck, J. R., Roellig, T. L., van Cleve, J., et al. 2004, *ApJS*, 154, 18
- Houck, J. R., Weedman, D. W., Le Floc'h, E., and Hao, L. 2007, *ApJ*, 671, 323
- Huynh, M. T., Norris, R. P., Coppin, K. E. K., et al. 2013, *MNRAS*, 431, 88
- Inami, H., Armus, L., Charmandaris, V. et al. 2013, *ApJ*, 777, 156
- Janssen, A. W., Christopher, N., Sturm, E. et al. 2016, *arxiv1604.00185*
- Laurent, O., Mirabel, R. F., Charmandaris, V., et al. 2000, *A&A*, 359, 887
- Lebouteiller, V., Bernard-Salas, J., Sloan, G. C., and Barry, D. J. 2010, *PASP*, 122, 231
- Lebouteiller, V., Barry, D. J., Bernard-Salas, H., Sloan, G. C., Houck, J. R., and Weedman, D. W. 2010, *ApJS*, 196, 8
- Lebouteiller, V., Cormier, D., Madden, S. C., et al. 2012, *A&A*, 548, A91
- Lebouteiller, V., Barry, D. J., Goes, C. et al. 2015, *ApJS*, 218, 21
- Luhman, M. L., Satyapal, S., Fischer, J., et al. 2003, *ApJ*, 594, 758
- Malhotra, S., Helou, G., Stacey, G., et al. 1997, *ApJ*, 491, L27

- Malhotra, S., Kaufman, M. J., Hollenbach, D., et al. 2001, *ApJ*, 561, 766
- Meijerink, R., Spaans, M., and Israel, F. P. 2007, *A&A*, 461, 793
- Nelson, C. H. and Whittle, M. 1996, *ApJ*, 465, 96
- Nikola, T., Genzel, R., Herrmann, F., et al. 1998, *ApJ*, 504, 749
- Petric, A. O., Armus, L., Howell, J., et al. 2011, *ApJ*, 730, 28
- Pilbratt, G. L., Riedinger, J. R., Passvogel, T., et al. 2010, *A&A*, 518, L1
- Poglitsch, A., Waelkens, C., Geis, N., et al. 2010, *A&A*, 518, L2
- Pottasch, S. and Bernard-Salas, J. 2010, *A&A*, 517, 95
- Sargsyan, L., Weedman, D., Lebouteiller, V., Houck, J., Barry, D., Hovhannisyan, A., and Mickaelian, A. 2011, *ApJ*, 730, 19
- Sargsyan, L., Lebouteiller, V., Weedman, D., et al. 2012, *ApJ*, 755, 171
- Sargsyan, L., Samsonyan, A., Lebouteiller, V. et al. 2014, *ApJ*, 790, 15
- Smith, M. G. and Weedman, D. W. 1972, *ApJ*, 172, 307
- Spoon, H. W. W., Farrah, D., Lebouteiller, V., et al. 2013, *ApJ*, 775, 127
- Stacey, G., Geis, N., Genzel, R., et al. 1991, *ApJ*, 373, 423
- Stierwalt, S., Armus, L., Surace, J. A., et al. 2013, *ApJS*, 206, 1
- Tielens, A. G. G. M., and Hollenbach, D. 1985, *ApJ*, 291, 722
- Tommasin, S., Spinoglio, L., Malkan, M., and Fazio, G. 2010, *ApJ*, 709, 1257
- Veilleux, S., Rupke, D. S. N., Kim, D.-C., et al. 2009, *ApJS*, 182, 628
- Wang, R., Wagg, J., Carilli, C.L., et al. 2013, *ApJ*, 773, 44
- Weedman, D.W. and Houck, J.R. 2009, *ApJ*, 693, 370
- Werner, M. W., Roellig, T. L., Low, F. J., et al. 2004, *ApJS*, 154, 1
- Whittle, M. 1985, *MNRAS*, 213, 1
- Wright, E. L. 2006, *PASP*, 118, 1711
- Wu, Y., Helou, G. , Armus, L., et al. 2010, *ApJ*, 723, 895

TABLE 1
NEON AND [C II] LINE WIDTHS AND REDSHIFTS

No.	Herschel AOR	Name ^a	R.A. ^b	Dec. ^b	z^c [C II]	$dV[\text{Ne II}]^d$ km s ⁻¹	$dV[\text{Ne III}]^e$ km s ⁻¹	FWHM ^f [C II]	FWHM ^g [Ne II]	FWHM ^h [Ne III]	EW ⁱ μm	Neon ^j ratio	spax ^k	size ^l kpc	ref ⁿ
1	1342235847	Mkn334	00:03:09.62	21:57:36.60	0.022151	31	98	249	328	460	0.25	0.41	3.3	4.15	9
2	1342235658	NGC0023	00:09:53.36	25:55:27.70	0.015119	33	-56	386	240	207	0.58	0.16	3.3	2.86	10
3	1342199416	NGC34/Mkn938	00:11:06.55	-12:06:26.30	0.019418	52	73	364	448	450	0.45	0.14	3.3	3.65	6
4	1342235846	MCG-02-01-051/2	00:18:50.90	-10:22:36.70	0.027243	13	34	197	321	300	0.72	0.22	3.3	5.08	9
5	1342237570	IRAS00188-0856	00:21:26.48	-08:39:27.10	0.128447	-12	92	310	328	959 ^P	0.07	0.15	3.3	21.30	8
6	1342237574	IRAS00199-7426	00:22:07.01	-74:09:41.70	0.096064	13	23	269	339	299	0.35	0.09	3.3	16.52	9
7	1342199236	Haro11	00:36:52.53	-33:33:18.60	0.020717	50	38	157	249	236	0.15	3.65	3.3	3.88	5
8	1342247543	NGC185	00:38:57.97	48:20:14.60	-0.000605	...	...	47	...	...	0.12	...	3.3	(0.03)	2
9	1342232583	E12-G21	00:40:46.16	-79:14:24.30	0.032967	42	42	333	381	276	0.28	0.49	3.3	6.10	9
10	1342238351	IRAS00397-1312	00:42:15.50	-12:56:03.50	...	...	...	452	316	316	0.03	0.5	3.3	37.67	8
11	1342238142	NGC0232a	00:42:45.83	-23:33:41.00	0.022890	53	107	449	210	373	0.55	0.16	3.3	4.28	10
12	1342238144	NGC0232b	00:42:52.82	-23:32:27.80	0.022302	71	40	499	520	477	0.16	0.9	3.4	4.18	10
13	1342199415	NGC253	00:47:33.12	-25:17:17.60	0.000773	41	53	67	342	352	0.34	0.07	3.4	(0.14)	1
14	1342238141	IRASF00456-2904SW	00:48:06.75	-28:48:18.60	0.109910	23	42	167	219	289	0.43	0.19	3.3	18.62	9
15	1342262037	IZw1	00:53:34.94	12:41:36.20	0.061170	140 ^q	...	310	331	...	<0.003	...	3.3	10.94	17
16	1342193211	MCG+12-02-001	00:54:03.88	73:05:05.90	0.016092	53	50	106	260	272	0.65	0.35	2.3	3.03	6
17	1342236877	NGC0317B	00:57:40.41	43:47:32.50	0.016949	3	28	399	364	302	0.57	0.09	3.3	3.20	10
18	1342262033	ESO541-IG12	01:02:17.55	-19:40:08.70	0.056695	-20	23	340	172	395	...	3.28	3.3	10.20	17
19	1342238371	IRAS01003-2238	01:02:49.94	-22:21:57.30	0.117807	-18	21	100	204	361	<0.043	0.31	3.3	19.77	8
20	1342247782	3C31	01:07:24.94	32:24:45.20	0.016941	55	2	699 ⁿ	607	544	0.11	...	3.3	3.19	16
21	1342212532	IC1623	01:07:47.57	-17:30:24.40	0.020201	161 ^q	182 ^r	264	255	264	0.3	0.89	2.3	3.79	6
22	1342238385	MCG-03-04-014	01:10:08.96	-16:51:09.80	0.035210	14	13	347	405	289	0.67	0.1	3.3	6.50	9
23	1342232616	NGC0454	01:14:24.90	-55:23:52.00	0.012129	114	98	202	427	348	<0.008	1.58	3.3	2.30	9
24	1342234998	ESO244-G012	01:18:08.31	-44:27:43.40	0.022489	-24	-10	199	350	365	0.66	0.23	3.3	4.21	9
25	1342238602	CGCG436-030	01:20:02.72	14:21:42.90	0.031363	82	103	220	250	215	0.5	0.27	3.3	5.81	10
26	1342238600	ESO353-G020	01:34:51.26	-36:08:14.40	0.016083	7	16	396	474	438	0.54	0.11	3.3	3.03	9
27	1342222218	NGC625-1-1	01:35:06.68	-41:26:13.00	0.001386	119	124	100	197	100	...	6.53	3.4	(0.17)	1
28	1342238556	ESO297-G011b	01:36:23.39	-37:19:18.10	0.017190	-5	-28	140	187	284	0.52	0.14	3.3	3.24	10
29	1342238558	ESO297-G011a	01:36:24.14	-37:20:25.90	...	...	...	149	231	231	0.7	0.29	3.5	3.26	10
30	1342238601	IRASF01364-1042	01:38:52.92	-10:27:11.40	0.048408	8	-11	384	395	488	0.39	0.22	3.3	8.79	9
31	1342238637	IRASF01417+1651	01:44:30.50	17:06:05.00	0.027452	-7	108	270	462	559	0.37	0.17	3.3	5.11	10
32	1342238392	NGC0695	01:51:14.24	22:34:56.50	0.032434	28	45	181	405	333	0.65	0.17	3.3	6.01	10
33	1342237475	UGC01385	01:54:53.82	36:55:04.30	0.018487	-7	-14	49	279	229	0.64	0.12	3.3	3.47	9
34	1342238723	Mkn1014	01:59:50.23	00:23:40.50	0.163097	-8	40	226	543	911	0.04	1.61	3.3	26.04	8
35	1342238364	NGC0788	02:01:06.45	-06:48:55.90	0.013539	-19	-24	152	230	265	<0.003	2.24	3.3	2.56	9
36	1342239376	IRAS02054+0835	02:08:06.90	08:50:04.30	...	...	...	447	...	...	0.02	...	3.3	45.67	9
37	1342238923	NGC0838c	02:09:24.61	-10:08:09.40	0.013568	78	28	345	528	300	0.45	0.21	3.3	2.56	10
38	1342238919	NGC0838a	02:09:38.60	-10:08:46.90	0.012798	48	47	205	206	73	0.74	0.28	3.3	2.42	10
39	1342238921	NGC0838b	02:09:42.75	-10:11:01.90	0.012922	-41	21	173	318	238	0.5	0.27	3.3	2.45	10
40	1342238724	NGC0828	02:10:09.53	39:11:24.70	0.017750	45	50	425	444	339	0.61	0.1	3.3	3.34	10
41	1342238950	IC0214	02:14:05.56	05:10:23.70	0.030017	-3	53	192	361	348	0.64	0.67	3.3	5.57	10
42	1342238636	Mkn590	02:14:33.56	-00:46:00.10	0.026591	-23	-4	218	835 ^o	629	<0.007	0.79	3.3	4.96	9
43	1342238956	NGC0877b	02:17:53.26	14:31:18.40	0.012795	-65 ^q	-48	118	365	338	0.46	0.18	3.4	2.42	10
44	1342238952	NGC0877a	02:17:59.68	14:32:38.20	0.013317	50	77	100	434	100	0.35	0.28	4.1	2.52	10
45	1342238422	MCG+05-06-36b	02:23:18.92	32:11:18.40	0.033639	24	7	246	410	487	0.54	0.07	3.5	6.22	10
46	1342238420	MCG+05-06-36a	02:23:21.99	32:11:49.50	0.034079	109	-23	523 ⁿ	445	587	0.48	0.2	3.3	6.29	10
47	1342239504	UGC01845	02:24:07.97	47:58:11.90	0.015702	56	105	400	365	313	0.58	0.1	3.3	2.96	9
48	1342238946	NGC0958	02:30:42.84	-02:56:20.40	0.019294	88	-7	250	415	100	0.29	0.32	4.4	3.63	10
49	1342239370	IC1816	02:31:51.00	-36:40:19.40	0.016937	-13	-2	125	440	502	0.03	1.27	3.3	3.19	9
50	1342239500	NGC0973	02:34:20.11	32:30:20.20	0.016215	-3	44	300	434	391	<0.002	1.21	3.3	3.05	9
51	1342238958	NGC0992	02:37:25.46	21:06:02.80	0.013630	58	35	240	142	121	0.72	0.24	3.3	2.57	10
52	1342191154	NGC1068	02:42:40.71	-00:00:47.80	0.003748	88	110	200	796 ^o	884 ^P	...	12.55	3.4	(0.57)	1
53	1342203121	NGC1068-1	02:42:40.71	-00:00:47.80	0.003721	80	102	204	796 ^o	884 ^P	...	12.55	3.3	(0.57)	1
54	1342203122	NGC1068-1	02:42:40.71	-00:00:47.80	0.003717	79	101	204	796 ^o	884 ^P	...	12.55	3.3	(0.57)	1
55	1342216210	NGC1056	02:42:48.47	28:34:24.70	0.005203	-107 ^q	-56	187	302	269	...	0.35	3.3	(1.22)	14
56	1342216212	NGC1056	02:42:48.47	28:34:24.70	0.005460	-30	21	68	302	269	...	0.35	3.4	(1.22)	14
57	1342238425	UGC02238	02:46:17.49	13:05:44.40	0.021696	15	15	342	249	227	0.63	0.17	3.3	4.07	10

TABLE 1 — *Continued*

No.	Herschel AOR	Name ^a	R.A. ^b	Dec. ^b	z ^c [C II]	dV[Ne II] ^d km s ⁻¹	dV[Ne III] ^e km s ⁻¹	FWHM ^f [C II]	FWHM ^g [Ne II]	FWHM ^h [Ne III]	EW ⁱ μm	Neon ^j ratio	spax ^k	size ^l kpc	ref ⁿ
58	1342239499	IRASF02437+2122	02:46:39.15	21:35:10.30	...	...	...	...	342	387	0.15	0.09	3.5	4.36	9
59	1342239497	UGC02369	02:54:01.79	14:58:25.00	0.031632	122	...	196	235	...	0.57	...	3.3	5.85	9
60	1342214034	NGC1140-1	02:54:33.58	-10:01:39.90	0.004961	19	-11	100	136	100	0.57	3.58	2.4	(0.83)	1
61	1342238915	Mkn1066	02:59:58.59	36:49:14.30	0.012189	7	36	202	331	465	0.25	0.46	3.3	2.31	9
62	1342239496	NGC1222	03:08:56.74	-02:57:18.50	0.008245	44	68	67	117	171	0.58	1.47	3.3	1.57	11
63	1342238960	UGC02608	03:15:01.47	42:02:08.60	0.023531	25	73	260	260	395	0.2	1.19	3.3	4.40	10
64	1342238963	IRAS03158+4227	03:19:12.60	42:38:28.00	0.134629	...	...	220	...	...	...	2.47	3.3	22.18	8
65	1342191351	NGC1275	03:19:48.16	41:30:42.10	0.017569	6	35	280	732 ^o	727	0.02	0.46	3.3	3.31	1
66	1342214362	NGC1275	03:19:48.16	41:30:42.10	0.017526	-7	22	287	732 ^o	737	<0.008	0.46	1.1	3.30	15
67	1342238940	IRASF03217+4022	03:25:05.37	40:33:32.20	0.023439	132 ^q	118	349	293	258	0.55	0.09	3.3	4.38	9
68	1342239752	Mkn609	03:25:25.34	-06:08:38.70	0.034600	35	51	100	273	327	0.32	0.29	3.3	6.39	9
69	1342191294	NGC1365	03:33:36.37	-36:08:25.50	0.005530	42	76	289	260	265	0.13	0.7	3.3	(0.83)	1
70	1342192138	NGC1386	03:36:46.20	-35:59:57.00	0.002536	-132 ^q	-128 ^f	142	271	365	<0.001	2.87	3.4	(0.74)	1
71	1342214221	SBS0335-052	03:37:44.00	-05:02:38.00	0.013585	...	86	100	...	92	<0.006	39.16	3.3	2.57	1
72	1342238916	IRASF03359+1523	03:38:47.14	15:32:53.80	0.036012	150 ^q	140	246	319	364	0.36	0.3	2.3	6.64	9
73	1342238943	IRASF03450+0055	03:47:40.18	01:05:14.00	...	...	...	...	325	600	<0.001	1.84	3.2	5.75	9
74	1342238967	CGCG465-012b	03:54:07.67	15:59:24.30	0.022214	92	67	134	271	258	0.67	0.13	3.5	4.16	10
75	1342238965	CGCG465-012a	03:54:15.95	15:55:43.40	0.022278	97	59	100	265	248	0.6	0.09	3.3	4.17	10
76	1342223119	IRAS03538-6432	03:54:25.21	-64:23:44.70	...	...	...	...	498	577	0.08	0.26	3.1	41.61	9
77	1342239746	IRAS03521+0028	03:54:42.15	00:37:02.00	0.151785	9	-9	587 ⁿ	425	604	0.25	0.39	3.3	24.54	8
78	1342243655	IRAS03582+6012	04:02:32.99	60:20:41.80	0.029984	42	84	183	78	100	0.64	0.48	3.3	5.57	10
79	1342239509	IRAS04103-2838	04:12:19.53	-28:30:24.40	0.117712	41	94	280	442	502	0.18	0.76	3.3	19.76	9
80	1342239505	UGC02982	04:12:22.68	05:32:49.10	0.017841	-5	51	285	189	128	0.68	0.14	3.3	3.36	10
81	1342226902	IRAS04114-5117	04:12:44.92	-51:09:34.20	0.125096	-122 ^q	128	100	564	1023 ^p	0.34	0.22	3.3	20.82	9
82	1342238379	ESO420-G013	04:13:49.70	-32:00:25.30	0.012080	23	30	57	299	255	0.3	0.22	3.3	2.29	9
83	1342239480	NGC1572	04:22:42.81	-40:36:03.10	0.020504	-2	31	366	404	415	0.46	0.08	3.3	3.85	10
84	1342239696	IRAS04271+3849	04:30:33.09	38:55:47.80	0.019178	8	10	369	332	294	0.61	0.1	3.3	3.61	10
85	1342225795	3C120	04:33:11.10	05:21:15.60	0.033501	143 ^q	133	289	285	302	<0.001	3.73	3.4	6.19	9
86	1342190367	NGC1614	04:33:59.85	-08:34:44.00	0.015835	68	7	245	346	381	0.61	0.96	3.3	2.98	6
87	1342239698	UGC03094	04:35:33.83	19:10:18.20	0.024673	-36	-4	339	296	289	0.43	0.33	3.3	4.61	10
88	1342238378	ESO203-IG001	04:46:49.55	-48:33:30.60	0.052950	37	185 ^f	281	442	100	0.03	0.22	3.3	9.56	9
89	1342239733	MCG-05-12-006	04:52:04.96	-32:59:26.00	0.018472	-4	-15	198	297	304	0.53	0.09	3.3	3.47	9
90	1342219440	NGC1808	05:07:42.32	-37:30:45.70	0.003385	74	45	267	350	441	0.58	0.36	3.3	(0.45)	1
91	1342270686	NGC1808	05:07:42.32	-37:30:45.70	0.003386	90	61	267	350	441	0.58	0.1	3.3	(0.45)	3
92	1342239765	NGC1797	05:07:44.84	-08:01:08.70	0.015021	17	-16	256	178	242	0.62	0.11	3.3	2.84	10
93	1342243518	CGCG468-002b	05:08:19.71	17:21:47.80	0.017603	-23	12	360	270	334	0.12	0.96	3.3	3.31	10
94	1342243520	CGCG468-002a	05:08:21.21	17:22:08.00	0.016725	53	38	202	228	198	0.54	0.11	3.3	3.16	10
95	1342239694	IRAS05083+2441	05:11:25.88	24:45:18.20	0.023535	59	74	262	254	78	0.72	0.32	3.3	4.40	10
96	1342226750	Akn120	05:16:11.42	-00:08:59.40	0.032865	11	49	297	349	424	<0.001	1.37	3.3	6.08	9
97	1342219853	VII Zw31	05:16:46.39	79:40:12.90	0.054229	-4	-26	113	217	227	0.64	0.11	3.4	9.80	9
98	1342243529	IRAS05129+5128	05:16:55.96	51:31:56.90	0.027664	16	23	133	254	242	0.54	0.94	3.3	5.15	10
99	1342219442	IRAS05189-2524	05:21:01.41	-25:21:45.50	0.042708	-6	419 ^f	181	345	754	0.02	0.93	3.3	7.81	1
100	1342227348	IRASF05187-1017	05:21:06.54	-10:14:46.70	0.028617	-38	-6	348	395	388	0.53	0.17	3.5	5.32	9
101	1342230077	IRAS05223+1908	05:25:16.69	19:10:47.80	...	...	...	...	...	...	<0.01	...	3.3	5.50	10
102	1342222085	30dorEstrip-1	05:38:34.58	-69:05:57.50	0.000912	27	56	45	212	264	<0.008	15.23	5.3	(0.002)	1
103	1342243527	MCG+08-11-002	05:40:43.70	49:41:41.60	0.019183	0	-19	458	548	438	0.56	0.09	3.3	3.61	10
104	1342230951	NGC1961	05:42:04.80	69:22:43.00	0.013053	-4	-13	280	568	618	0.24	0.31	4.3	2.47	10
105	1342243531	UGC03351	05:45:48.03	58:42:03.60	0.014914	-31	37	276	352	339	0.53	0.12	3.3	2.82	10
106	1342243522	IRAS05442+1732	05:47:11.20	17:33:46.40	0.018485	-35	-12	249	312	274	0.66	0.33	3.3	3.47	10
107	1342239744	2MASXJ0558-3820	05:58:02.00	-38:20:04.70	0.034224	212 ^q	122	100	524	467	<0.001	1.69	3.3	6.32	9
108	1342239479	IRAS06035-7102	06:02:53.63	-71:03:11.90	0.079450	-65 ^q	23	236	415	522	<0.076	0.27	3.3	13.93	8
109	1342230910	IRASF06076-2139	06:09:45.74	-21:40:24.50	0.037644	20	37	142	359	464	0.33	0.07	3.3	6.93	9
110	1342230944	UGC03410b	06:13:58.59	80:28:35.50	0.012585	-54	-32	112	268	311	0.56	0.13	3.3	2.38	10
111	1342230997	UGC03410a	06:14:30.35	80:27:00.20	0.013040	36	18	182	318	422	0.63	0.11	3.3	2.47	10
112	1342219851	Mkn334	06:15:36.36	71:02:15.10	0.013450	-51	-58 ^f	332	729 ^o	748	<0.002	1.83	3.3	2.54	1
113	1342193210	NGC2146-nuc	06:18:37.71	78:21:25.30	0.002792	-69 ^q	-45	143	293	205	0.67	0.64	2.3	(1.00)	6
114	1342230961	IRAS06206-6315	06:21:00.80	-63:17:23.20	0.092098	54	181 ^f	385	523	608	<0.107	0.52	3.3	15.91	8
115	1342231278	IRAS06301-7934	06:26:42.20	-79:36:30.40	0.156284	-60	-343 ^f	271	437	495	0.15	0.38	3.3	25.14	9
116	1342226880	ESO255-IG007a	06:27:21.70	-47:10:36.20	0.039435	-10	-7	210	395	349	0.62	0.22	3.3	7.23	10

TABLE 1 — *Continued*

No.	Herschel AOR	Name ^a	R.A. ^b	Dec. ^b	z ^c [C II]	dV[Ne II] ^d km s ⁻¹	dV[Ne III] ^e km s ⁻¹	FWHM ^f [C II]	FWHM ^g [Ne II]	FWHM ^h [Ne III]	EW ⁱ μm	Neon ^j ratio	spax ^k	size ^l kpc	ref ⁿ
117	1342226190	ESO255-IG007b	06:27:23.09	-47:11:02.60	0.039165	-44	-47	220	172	100	0.65	0.6	3.3	7.19	10
118	1342230958	ESO557-G002b	06:31:45.71	-17:38:44.90	0.021277	3	-8	157	128	196	0.7	0.49	3.5	3.98	10
119	1342230956	ESO557-G002a	06:31:47.20	-17:37:16.60	0.021308	38	61	67	165	219	0.6	0.15	3.3	3.99	10
120	1342238377	IRAS06361-6217	06:36:35.85	-62:20:32.60	0.160096	113	64	270	247	361	0.05	0.25	3.3	25.65	9
121	1342230996	NGC2273	06:50:08.72	60:50:45.00	0.006183	-81 ^q	-42	276	202	289	0.13	0.57	3.3	(1.41)	9
122	1342230955	UGC03608	06:57:34.41	46:24:10.60	0.021647	85	45	149	174	207	0.53	0.14	3.3	4.06	9
123	1342231286	IRASF06592-6313	06:59:40.26	-63:17:52.40	0.023044	10	22	299	353	449	0.48	0.1	3.3	4.31	9
124	1342226876	IRASF07027-6011a	07:03:24.14	-60:15:21.90	0.031754	65	285 ^r	161	281	648	0.04	0.68	3.3	5.89	10
125	1342226878	IRASF07027-6011b	07:03:28.52	-60:16:43.70	0.031019	32	18	81	255	217	0.68	0.1	3.5	5.75	10
126	1342229686	NGC2342a	07:09:18.08	20:38:09.50	0.017579	-42	-19	247	202	240	0.67	0.15	3.3	3.31	10
127	1342207825	IRAS07251-0248	07:27:37.62	-02:54:54.80	0.087810	-9	-15	331	442	183	0.09	0.12	3.3	15.24	1
128	1342229716	NGC2388a	07:28:53.44	33:49:08.80	0.013723	1	11	203	257	236	0.53	0.08	3.3	2.60	10
129	1342230946	MCG+02-20-003	07:35:43.44	11:42:34.80	0.016580	8	25	209	239	289	0.17	0.11	3.3	3.13	10
130	1342243533	Mkn9	07:36:56.98	58:46:13.40	0.040100	-7	22	100	278	215	<0.142	1.06	3.3	7.36	9
131	1342243534	IRAS07598+6508	08:04:30.46	64:59:52.90	0.148697	95	1449 ^r	211	528	1044 ^p	0.01	0.39	3.3	24.12	9
132	1342229688	Mkn622	08:07:41.04	39:00:15.20	0.023510	148 ^q	97	148	414	821 ^p	0.24	0.52	3.3	4.39	9
133	1342220753	HS0822+3542	08:25:55.52	35:32:32.00	...	...	...	...	...	100	<0.13	...	3.3	0.47	5
134	1342230967	IRAS08311-2459	08:33:20.47	-25:09:33.10	0.100478	-53	-58 ^r	244	265	717	0.14	1.03	3.3	17.20	8
135	1342186306	He2-10	08:36:15.18	-26:24:33.90	...	...	...	...	...	189	0.61	1.23	3.4	(0.43)	1
136	1342221975	He2-10	08:36:15.18	-26:24:33.90	0.002864	4	12	60	189	121	0.61	1.23	3.3	(0.43)	1
137	1342225823	IRAS08355-4944	08:37:01.87	-49:54:30.00	0.025928	57	106	91	290	131	0.19	0.56	3.3	4.83	10
138	1342208904	NGC2623	08:38:24.08	25:45:16.90	0.018427	43	84	344	392	469	0.27	0.27	3.3	3.46	6
139	1342230964	ESO432-IG006b	08:44:27.21	-31:41:50.80	0.016292	-60	-5	325	214	287	0.64	0.12	3.5	3.08	10
140	1342230962	ESO432-IG006a	08:44:28.93	-31:41:30.30	0.016257	-21	13	394	437	463	0.41	0.14	3.3	3.07	10
141	1342228521	ESO60-IG016	08:52:32.07	-69:01:54.80	0.044863	13	45	192	144	287	0.07	0.75	3.3	8.18	9
142	1342208952	IRAS08572+3915	09:00:25.38	39:03:54.30	0.058238	-8	...	198	226	...	<0.03	...	3.3	10.46	1
143	1342231958	Mkn18	09:01:58.39	60:09:06.20	0.011332	63	50	236	170	192	0.27	0.58	3.3	2.15	9
144	1342209403	IRAS09022-3615	09:04:12.00	-36:27:01.00	0.059827	-32	-41	343	352	402	0.14	0.58	2.3	10.72	1
145	1342230935	UGC04881	09:15:55.10	44:19:55.00	0.039950	32	15	194	354	267	0.4	0.08	3.3	7.33	10
146	1342231718	MCG-01-24-012	09:20:46.25	-08:03:22.10	0.019814	27	28	268	328	371	<0.005	0.96	3.3	3.72	9
147	1342231715	Mkn705	09:26:03.29	12:44:03.60	0.028791	-7	43	285	183	396	0.02	1.03	3.3	5.35	9
148	1342220973	I Zw18	09:34:02.03	55:14:28.10	0.002585	-8	27	66	100	100	<0.09	8.49	2.3	(0.86)	1
149	1342208949	UGC5101	09:35:51.65	61:21:11.30	0.039277	71	87	498	477	706	0.18	0.37	3.3	7.21	1
150	1342231716	IC0563	09:46:20.30	03:02:44.70	0.020094	16	-30	225	392	410	0.6	0.11	3.3	3.77	10
151	1342231731	IRASF09437+0317	09:46:21.10	03:04:16.30	0.020027	19	29	195	416	321	0.57	0.09	3.3	3.76	10
152	1342221391	NGC3079	10:01:57.80	55:40:47.10	0.003843	75	99	452	570	633	0.4	0.25	3.3	(0.88)	1
153	1342270667	Mkn25	10:03:51.90	59:26:10.00	0.010103	41	48	100	136	27	0.46	0.46	3.3	1.92	3
154	1342231725	NGC3110a	10:04:02.11	-06:28:29.20	0.016982	28	32	280	338	200	0.64	0.08	3.3	3.20	10
155	1342254278	3C236	10:06:01.74	34:54:10.40	0.099224	-11	-25	491	1298 ^o	541	...	0.4	3.3	17.00	4
156	1342235705	IRASF10038-3338	10:06:04.65	-33:53:06.10	0.034215	29	-26	90	180	181	<0.002	0.29	3.3	6.31	9
157	1342231729	IRASF10173+0828	10:20:00.22	08:13:34.00	0.048963	-32	-384 ^r	294	364	100	0.35	0.24	3.3	8.89	10
158	1342233703	NGC3221	10:22:20.26	21:34:22.30	0.013379	4	59	73	154	100	0.75	0.42	4.3	2.53	10
159	1342197796	NGC3227	10:23:30.60	19:51:54.00	0.003907	33	60	290	465	508	0.09	1.1	3.3	(0.99)	14
160	1342232315	IRAS10378+1109	10:40:29.17	10:53:17.70	0.136400	-78 ^q	28	373	477	295	<0.078	0.2	3.3	22.43	8
161	1342225718	ESO264-G036	10:43:07.51	-46:12:44.10	0.020642	139 ^q	144	271	404	217	0.44	0.12	3.3	3.87	10
162	1342221892	Har03	10:45:22.33	55:57:37.30	0.003205	51	73	100	170	100	0.51	2.95	3.3	(0.86)	5
163	1342232587	NGC3393	10:48:23.39	-25:09:42.80	0.012587	9	14	171	425	454	<0.003	3.89	3.3	2.38	9
164	1342209015	Mkn153	10:49:05.03	52:20:07.80	...	...	...	...	303	128	<0.07	6.41	3.4	1.52	5
165	1342225716	ESO264-G057	10:59:01.70	-43:26:25.20	0.017158	-28	11	142	279	207	0.6	0.07	3.3	3.23	10
166	1342207788	IRAS10565+2448	10:59:18.14	24:32:34.30	0.043080	7	8	139	242	292	0.38	0.12	3.3	7.88	1
167	1342223365	NGC3489	11:00:18.57	13:54:04.40	0.002331	-17	14	152	151	151	...	1.53	3.3	(0.41)	11
168	1342233712	MCG+07-23-019	11:03:53.20	40:50:57.00	0.035245	84	77	234	435	380	0.64	0.2	2.2	6.50	10
169	1342245980	NGC3516	11:06:47.49	72:34:06.90	0.009168	63	113	86	613	403	...	1.98	3.4	(2.55)	14
170	1342232612	IRAS11095-0238	11:12:03.34	-02:54:24.10	0.106358	30	17	208	322	276	<0.027	0.38	3.3	18.08	8
171	1342232307	IRAS11119+3257	11:14:38.88	32:41:33.10	...	...	...	...	360	937 ^p	0.01	0.58	3.3	29.35	9
172	1342223770	CGCG011-076	11:21:12.26	-02:59:03.50	0.024900	62	43	198	490	476	0.32	0.12	3.3	4.43	10
173	1342223753	IC2810a	11:25:45.05	14:40:35.90	0.034607	178 ^q	189 ^r	229	393	375	0.6	0.11	3.4	6.39	10
174	1342235704	ESO319-G022	11:27:54.18	-41:36:51.70	0.016339	5	28	171	302	367	0.3	0.07	3.3	3.09	9
175	1342199289	VIIZw31403	11:27:59.90	78:59:39.00	-0.000289	...	...	46	...	100	<0.16	13.32	4.2	(0.05)	5

TABLE 1 — *Continued*

No.	Herschel AOR	Name ^a	R.A. ^b	Dec. ^b	z ^c [C II]	$dV[\text{Ne II}]^d$ km s ⁻¹	$dV[\text{Ne III}]^e$ km s ⁻¹	FWHM ^f [C II]	FWHM ^g [Ne II]	FWHM ^h [Ne III]	EW ⁱ μm	Neon ^j ratio	spax ^k	size ^l kpc	ref ⁿ
176	1342208906	Arp299	11:28:31.19	58:33:49.00	0.010944	101	86	202	268	248	...	0.48	5.4	2.07	1
177	1342199419	NGC3690/Arp299	11:28:31.32	58:33:42.50	0.010635	123	163 ^r	151	100	340	...	0.34	3.3	2.02	1
178	1342222070	Mkn1450	11:38:35.62	57:52:27.20	0.003202	-17	7	100	159	142	0.26	10.66	3.3	(0.61)	5
179	1342212225	NGC3783	11:39:01.78	-37:44:18.70	0.009825	82	106	59	245	229	<0.001	1.47	3.3	(2.18)	1
180	1342222201	UM448	11:42:12.30	00:20:03.00	0.018460	10	-6	145	274	242	0.45	2.27	3.3	3.47	5
181	1342245394	NGC3894	11:48:50.36	59:24:56.40	...	...	...	...	787 ^o	626	0.02	0.98	3.3	2.04	16
182	1342222205	UM461	11:51:33.04	-02:22:22.80	0.003610	...	32	120	...	100	<0.05	18.54	3.3	(0.69)	5
183	1342197912	NGC3982	11:56:28.10	55:07:30.60	0.003704	-44	-29	79	279	244	...	0.99	3.3	(0.99)	14
184	1342210822	Antennae/NGC4038	12:01:54.75	-18:52:51.10	0.005265	115	185 ^r	95	242	248	...	0.95	3.3	(0.97)	1
185	1342199228	SBS1159+545	12:02:02.44	54:15:50.10	...	...	...	...	...	...	<0.04	...	3.3	2.27	5
186	1342197812	NGC4051	12:03:09.61	44:31:52.80	0.002354	4	39	66	244	336	0.05	1.12	3.3	(0.66)	1
187	1342208880	NGC4051	12:03:09.61	44:31:52.80	0.002317	-7	27	50	244	336	0.05	1.12	3.3	(0.66)	14
188	1342223403	IRAS12018+1941	12:04:24.53	19:25:09.80	0.167849	34	-11	100	287	389	0.11	0.15	3.3	26.67	9
189	1342223404	UGC07064	12:04:43.34	31:10:38.20	0.025079	20	30	100	244	246	0.08	0.98	3.3	4.68	9
190	1342225714	ESO440-IG058	12:06:51.87	-31:56:59.20	0.022994	19	23	358	369	395	0.66	0.13	3.3	4.30	10
191	1342234994	IRAS12071-0444	12:09:45.12	-05:01:13.90	0.128861	1	311 ^r	251	305	859 ^p	0.05	0.96	3.3	21.36	8
192	1342198305	NGC4151	12:10:32.58	39:24:20.60	0.003396	80	104	196	350	325	<0.011	1.92	3.3	(0.64)	1
193	1342223773	NGC4150	12:10:33.65	30:24:05.50	0.000838	36	33	149	324	348	...	0.75	3.3	(0.58)	11
194	1342210832	IRASF12112+0305	12:13:46.05	02:48:41.30	0.072923	64	139	390	443	308	0.3	0.25	3.3	12.87	1
195	1342225793	ESO267-G030b	12:13:52.28	-47:16:25.40	0.018257	40	31	200	201	361	0.68	0.14	3.3	3.43	10
196	1342233265	NGC4194	12:14:09.64	54:31:34.60	0.008589	99	85	194	334	267	0.55	0.52	3.3	(1.78)	10
197	1342225812	ESO267-G030a	12:14:12.81	-47:13:42.50	0.018111	29	24	355	506	355	0.5	0.24	3.3	3.41	10
198	1342226873	IRAS12116-5615	12:14:22.08	-56:32:32.70	0.027267	13	22	67	239	282	0.36	0.12	3.3	5.08	10
199	1342199408	Tol1214-277	12:17:17.09	-28:02:32.70	...	...	...	...	...	153	<0.16	7.43	3.3	4.84	5
200	1342234057	IRASF12224-0624	12:25:03.91	-06:40:52.60	0.026433	9	...	208	309	...	0.07	...	3.3	4.92	10
201	1342197909	NGC4388	12:25:46.75	12:39:43.50	0.008517	91	68	208	276	316	0.03	1.33	3.3	(0.92)	14
202	1342199423	Mkn209	12:26:16.02	48:29:36.60	0.001039	87	51	100	202	272	<0.18	43.56	3.3	(0.26)	5
203	1342210830	NGC4418	12:26:54.66	-00:52:40.30	0.007147	...	...	109	...	...	<0.02	...	3.3	(1.15)	1
204	1342202109	NGC4418-1	12:26:54.66	-00:52:40.30	0.007083	...	...	93	...	...	<0.02	...	3.3	(1.15)	1
205	1342223803	NGC4435	12:27:40.49	13:04:44.20	0.002738	-31	-30	402	398	62	0.23	0.2	3.3	(0.78)	11
206	1342234949	NGC4459	12:29:00.03	13:58:42.80	0.004103	63	-1	362	240	100	...	1.1	3.3	(0.75)	11
207	1342235707	3C273	12:29:06.67	02:03:08.10	...	...	...	...	538	736	<0.002	2.61	3.3	25.42	8
208	1342234992	NGC4472	12:29:46.78	08:00:01.90	...	...	...	...	700 ^o	523	<0.002	0.54	3.3	0.63	12
209	1342225720	NGC4507	12:35:36.55	-39:54:33.30	0.011874	10	52	287	383	392	<0.002	0.96	3.3	2.25	9
210	1342198302	NGC4593	12:39:39.43	-05:20:39.30	0.008407	66	55	247	180	171	<0.006	1.21	3.3	(1.58)	1
211	1342212224	TOL1238-364	12:40:52.86	-36:45:21.10	0.011083	50	24	100	223	388	...	0.7	3.4	2.07	14
212	1342212222	TOL1238-364	12:40:52.86	-36:45:21.10	0.011083	50	24	100	223	388	...	0.7	3.4	2.07	14
213	1342236279	PG1244+026	12:46:35.24	02:22:08.70	0.048578	-45	54	100	255	384	0.02	0.58	3.3	8.83	9
214	1342237582	IRAS12514+1027	12:54:00.82	10:11:12.40	...	...	...	...	758 ^o	...	<0.009	...	3.3	43.26	9
215	1342186811	Mkn231	12:56:14.29	56:52:25.10	0.042180	34	...	153	193	...	0.01	...	3.3	7.71	1
216	1342223759	NGC4922	13:01:25.26	29:18:49.60	0.023740	31	40	167	223	384	0.16	0.27	3.3	4.43	10
217	1342225721	CGCG043-099	13:01:50.80	04:20:00.00	0.037912	118	124	278	359	313	0.55	0.1	2.2	6.97	10
218	1342225747	MCG-02-33-098	13:02:19.66	-15:46:04.20	0.015976	43	14	118	230	360	0.55	0.23	3.3	3.01	10
219	1342225742	ESO507-G070	13:02:52.42	-23:55:17.80	0.021534	5	-9	500	517	443	0.56	0.15	3.3	4.04	9
220	1342225749	NGC4941	13:04:13.06	-05:33:05.80	0.003718	-56	-18	156	212	276	<0.041	2.03	3.3	(0.78)	9
221	1342225811	ESO323-G077	13:06:26.13	-40:24:52.50	0.015786	-6	21	400	482	548	0.03	0.59	3.3	2.98	9
222	1342226187	IRAS13052-5711	13:08:18.73	-57:27:30.30	0.021851	56	56	399	371	321	0.61	0.17	3.3	4.10	10
223	1342234990	IC0860	13:15:03.53	24:37:07.90	0.012973	4	-26	218	318	100	0.43	0.14	3.3	2.45	10
224	1342214629	IRAS13120-5453	13:15:06.40	-55:09:23.30	0.031083	-38	31	340	288	411	0.45	0.16	3.3	5.76	1
225	1342232597	VV250b	13:15:30.69	62:07:45.80	...	...	...	...	555	610	0.76	0.31	3.5	5.71	10
226	1342232595	VV250a	13:15:34.96	62:07:29.20	0.031262	26	45	337	435	486	0.63	0.37	3.3	5.79	10
227	1342197801	Arp193	13:20:35.34	34:08:22.20	0.023412	65	27	391	317	292	0.62	0.17	3.3	4.38	6
228	1342225750	NGC5104	13:21:23.08	00:20:32.70	0.018611	102	109	495	365	462	0.51	0.11	3.3	3.50	10
229	1342225746	MCG-03-34-064	13:22:24.45	-16:43:42.40	...	...	...	...	816 ^o	801 ^p	<0.01	2.18	3.5	3.12	9
230	1342202589	CenA	13:25:27.62	-43:01:08.80	0.001838	54	62	122	520	428	<0.003	14.98	3.4	(0.17)	1
231	1342190371	NGC5135	13:25:43.97	-29:50:00.00	0.013717	81	131	57	302	346	0.39	0.46	3.3	2.60	6
232	1342190368	ESO173-G015	13:27:23.79	-57:29:21.80	0.010034	33	45	183	303	238	0.53	0.2	3.3	1.91	6
233	1342232550	IRASF13279+3401	13:30:15.23	33:46:29.40	...	...	...	...	154	...	0.04	...	3.4	4.44	9
234	1342237579	IC4280	13:32:53.40	-24:12:25.50	0.016255	-35	18	231	486	407	0.59	0.08	3.3	3.07	10

TABLE 1 — *Continued*

No.	Herschel AOR	Name ^a	R.A. ^b	Dec. ^b	z ^c [C II]	dV[Ne II] ^d km s ⁻¹	dV[Ne III] ^e km s ⁻¹	FWHM ^f [C II]	FWHM ^g [Ne II]	FWHM ^h [Ne III]	EW ⁱ μm	Neon ^j ratio	spax ^k	size ^l kpc	ref ⁿ
235	1342225810	M-6-30-15	13:35:53.78	-34:17:44.10	0.008006	40	70	106	208	100	<0.001	1.87	4.3	(1.52)	9
236	1342234948	IRAS13342+3932	13:36:24.07	39:17:30.10	0.179642	87	158	375	328	562	0.07	0.98	3.3	28.18	9
237	1342231427	IRAS13352+6402	13:36:51.15	63:47:04.70	...	...	...	...	352	948 ^P	0.08	0.26	3.3	34.95	9
238	1342236984	IRAS13349+2438	13:37:18.73	24:23:03.30	0.108287	...	-121 ^r	387	...	415	<0.001	...	3.3	18.37	9
239	1342231735	NGC5256	13:38:17.69	48:16:33.90	0.028120	-9	85	449	468	559	0.44	0.87	3.3	5.23	10
240	1342225569	NGC5257a	13:39:52.95	00:50:25.90	0.022732	...	...	215	...	...	0.63	...	3.2	4.25	10
241	1342225571	NGC5257b	13:39:57.72	00:49:53.00	0.022533	100	79	261	414	238	0.54	0.11	2.2	4.22	10
242	1342207802	Mkn273	13:44:42.12	55:53:13.10	0.037816	103	77	496	534	605	0.09	0.68	3.3	6.96	1
243	1342236886	IRAS13451+1232	13:47:33.36	12:17:24.20	0.121928	140 ^q	592 ^r	470	743 ^o	1801	<0.003	1.2	3.3	20.37	8
244	1342233263	UGC08739	13:49:13.94	35:15:26.20	0.016874	44	37	330	249	134	0.51	0.14	2.3	3.18	10
245	1342223809	IC4329A	13:49:19.29	-30:18:34.40	0.015966	72	90	450	521	589	<0.001	2.45	3.3	3.01	1
246	1342238373	ESO221-IG010	13:50:56.92	-49:03:18.80	0.010544	63	90	146	204	284	0.69	0.13	3.3	2.00	10
247	1342224240	NGC5331a	13:52:16.21	02:06:05.10	0.033202	41	71	480	512	520	0.61	0.12	3.3	6.14	10
248	1342225723	NGC5331b	13:52:16.43	02:06:30.90	0.033086	3	0	431	485	328	0.6	0.1	3.3	6.12	10
249	1342223133	NGC5347	13:53:17.83	33:29:27.00	0.008137	-1	13	100	90	346	0.03	1.16	3.3	(1.25)	9
250	1342236982	Mkn463	13:56:02.90	18:22:19.00	0.050817	68	22	219	667	555	<0.001	3.52	3.3	9.21	8
251	1342222240	NGC5394	13:58:33.65	37:27:12.60	0.011634	30	34	100	204	153	0.62	0.11	3.3	2.21	10
252	1342236271	IRAS14026+4341	14:04:38.72	43:27:07.30	...	...	...	...	353	...	0.01	...	3.4	43.73	9
253	1342247534	OQ208	14:07:00.39	28:27:14.70	0.076773	19	-62 ^r	474	488	776 ^P	0.02	0.58	3.3	13.50	4
254	1342237584	OQ208	14:07:00.39	28:27:14.70	...	...	...	...	488	776 ^P	0.02	0.58	3.5	13.47	9
255	1342191297	CircinusGalaxy	14:13:09.91	-65:20:20.50	0.001456	-1	7	156	245	173	0.04	3.4	3.3	(0.19)	1
256	1342213758	NGC5506	14:13:14.87	-03:12:27.00	0.006192	65	88	241	529	506	<0.007	1.52	3.3	(1.08)	1
257	1342199733	SBS1415+437	14:17:01.76	43:30:13.40	0.002093	...	...	100	...	...	<0.097	...	3.3	(0.62)	5
258	1342236985	NGC5548	14:17:59.53	25:08:12.40	0.017262	32	39	100	400	311	0.02	1	3.3	3.25	9
259	1342232331	Mkn1490	14:19:43.25	49:14:11.70	...	...	...	...	201	311	0.56	0.1	3.5	4.79	9
260	1342238159	PG1426+015	14:29:06.59	01:17:06.50	0.086587	14	36	287	551	572	<0.002	1.91	3.3	15.05	9
261	1342238163	NGC5653	14:30:10.44	31:12:55.80	0.011737	-68 ^q	-26	120	156	260	0.54	0.14	3.3	2.23	10
262	1342224242	IRAS14348-1447	14:37:38.27	-15:00:24.60	0.082551	-5	-32	286	454	299	<0.174	0.18	3.3	14.42	1
263	1342204338	IRAS14378-3651	14:40:58.90	-37:04:33.00	0.068118	124	124	148	341	231	0.39	0.08	3.3	12.10	1
264	1342223736	PG1440+356	14:42:07.46	35:26:22.90	0.077692	75	6	182	327	453	0.02	0.81	3.3	13.64	9
265	1342238133	NGC5728	14:42:23.93	-17:15:11.40	0.009492	60	16	351	514	668	0.08	2.34	3.3	(1.39)	9
266	1342238131	NGC5734a	14:45:09.04	-20:52:13.20	0.013214	-91 ^q	-138 ^r	105	535	411	0.47	0.15	3.5	2.50	10
267	1342238129	NGC5734b	14:45:11.02	-20:54:48.60	0.014030	74	41	141	423	281	0.49	0.19	3.3	2.66	10
268	1342248560	3C305	14:49:21.80	63:16:15.30	0.041809	59	6	802 ⁿ	843 ^o	918	...	1.96	3.3	7.65	4
269	1342236986	VV340b	14:57:00.32	24:36:24.60	0.032869	36	-7	100	247	100	0.58	0.12	3.3	6.08	10
270	1342234963	VV340a	14:57:00.70	24:37:05.80	0.034909	95	125	264	273	333	0.58	0.29	3.3	6.44	10
271	1342239742	IC4518ABa	14:57:41.18	-43:07:55.60	0.015817	-20	44	229	392	381	0.06	1.46	3.3	2.98	10
272	1342238134	NGC5793	14:59:24.76	-16:41:36.10	0.011596	43	55	401	361	351	0.39	0.2	3.3	2.20	9
273	1342238158	NGC5813	15:01:11.25	01:42:07.10	0.006891	83	-101 ^r	227	273	593	<0.016	1.06	3.3	(1.33)	12
274	1342236887	IRAS15001+1433	15:02:31.94	14:21:35.30	0.162125	60	65	378	488	675	0.14	0.36	3.3	25.92	9
275	1342238157	NGC5846	15:06:29.26	01:36:20.00	0.005721	-16	25	449	52	153	...	1.2	3.3	(1.23)	12
276	1342223161	VV705	15:18:06.34	42:44:36.70	0.040912	165 ^q	173 ^r	272	254	255	0.75	0.21	2.3	7.50	10
277	1342238152	IRAS15225+2350	15:24:43.94	23:40:10.20	0.138699	-29	344 ^r	100	450	870 ^P	0.05	0.31	3.3	22.74	9
278	1342238352	ESO099-G004	15:24:57.98	-63:07:29.40	0.029281	-49	-41	279	294	277	0.53	0.2	3.3	5.44	10
279	1342213752	IRAS15250+3609	15:26:59.40	35:58:37.50	0.055377	49	117	239	290	375	0.03	0.25	3.3	9.97	1
280	1342238153	NGC5936	15:30:00.85	12:59:22.10	0.013487	-6	7	100	208	164	0.62	0.08	3.3	2.55	10
281	1342191306	Arp220	15:34:57.24	23:30:11.70	0.018101	-8	4	470	369	486	0.17	0.14	3.3	3.40	1
282	1342238136	NGC5990	15:46:16.41	02:24:55.60	0.012594	-25	8	214	535	606	0.15	0.4	3.3	2.39	10
283	1342238135	IRAS15462-0450	15:48:56.80	-04:59:33.70	0.100358	-2	274 ^r	102	294	1296 ^P	0.05	0.51	3.3	17.18	8
284	1342253766	PKS1549-79	15:56:58.87	-79:14:04.30	0.152545	192 ^q	...	367	954 ^o	...	<0.008	...	3.3	24.64	4
285	1342237487	NGC6052	16:05:12.87	20:32:33.00	0.015809	52	53	80	214	169	0.69	0.76	3.3	2.98	10
286	1342234492	16052+5334	16:06:32.99	53:26:32.10	0.087864	-74 ^q	-33	636 ⁿ	387	327	...	0.13	3.3	15.25	7
287	1342238907	IRAS16090-0139	16:11:40.42	-01:47:05.80	0.133631	37	21	311	474	322	<0.057	0.17	3.3	22.04	8
288	1342227214	NGC6090	16:11:40.70	52:27:24.00	0.029573	-116 ^q	-16	55	139	276	0.73	0.27	3.3	5.50	10
289	1342222163	Mkn876	16:13:57.18	65:43:09.60	0.129276	-54	7	384	510	531	0.01	0.7	3.3	21.43	9
290	1342227799	IRAS16164-0746	16:19:11.79	-07:54:02.80	0.023400	14	35	374	467	458	0.42	0.3	3.2	4.37	9
291	1342238911	Mkn883	16:29:52.85	24:26:38.30	0.038311	7	68	296	291	473	0.26	0.65	3.3	7.04	9
292	1342238909	CGCG052-037	16:30:56.54	04:04:58.40	0.024785	15	25	198	204	200	0.62	0.08	3.3	4.63	9
293	1342232265	IRAS16334+4630	16:34:52.37	46:24:53.00	0.190795	15	62	263	452	646	0.3	0.24	3.3	29.58	9

TABLE 1 — *Continued*

No.	Herschel AOR	Name ^a	R.A. ^b	Dec. ^b	z ^c [C II]	dV[Ne II] ^d km s ⁻¹	dV[Ne III] ^e km s ⁻¹	FWHM ^f [C II]	FWHM ^g [Ne II]	FWHM ^h [Ne III]	EW ⁱ μm	Neon ^j ratio	spax ^k	size ^l kpc	ref ⁿ
294	1342238731	NGC6156	16:34:52.55	-60:37:08.00	0.011001	-6	51	146	299	377	0.37	0.29	3.3	2.08	10
295	1342231301	ESO069-IG006	16:38:11.85	-68:26:08.20	0.046332	60	31	497	491	609	0.64	0.14	3.3	8.44	9
296	1342240777	IRASF16399-0937N	16:42:40.11	-09:43:13.70	0.027296	15	1	248	554	725	0.38	0.31	3.3	5.09	9
297	1342250583	ESO453-G005	16:47:31.09	-29:21:21.60	0.021071	21	44	53	274	242	0.42	0.23	3.3	3.95	10
298	1342216623	NGC6240	16:52:58.89	02:24:03.40	0.024818	119	57	561 ⁿ	832 ^o	803 ^p	0.35	0.51	3.3	4.63	1
299	1342239382	IRASF16516-0948	16:54:23.72	-09:53:20.90	0.023099	-17	5	215	388	417	0.69	0.37	3.3	4.32	10
300	1342230134	NGC6286b/6285	16:58:24.01	58:57:21.50	0.019076	14	14	300	383	375	0.66	0.18	3.3	3.59	10
301	1342230132	NGC6286a	16:58:31.38	58:56:10.50	0.018497	45	65	286	322	315	0.59	0.1	3.3	3.48	10
302	1342238910	2MASSJ1659+1834	16:59:39.80	18:34:36.90	...	...	...	...	288	471	0.02	1.82	3.3	27.04	9
303	1342225993	PG1700+518	17:01:24.80	51:49:20.00	...	...	...	...	...	...	<0.009	...	3.3	40.76	9
304	1342223716	IRAS17044+6720	17:04:28.41	67:16:28.50	0.135376	21	127	219	477	757	0.04	0.4	3.3	22.28	9
305	1342232574	IRAS17068+4027	17:08:32.12	40:23:28.20	0.179438	113	107	375	435	392	<0.079	0.53	3.3	28.16	9
306	1342223740	IRASF17132+5313	17:14:20.00	53:10:30.00	0.051145	76	56	395	415	394	0.42	0.14	3.3	9.26	9
307	1342243510	IRASF17138-1017	17:16:35.60	-10:20:38.00	0.017492	60	127	199	346	262	0.68	0.11	3.3	3.29	10
308	1342229693	IRAS17208-0014	17:23:21.93	-00:17:00.40	0.042812	39	48	480	475	455	0.31	0.18	3.3	7.83	1
309	1342240168	ESO138-G027	17:26:43.35	-59:55:55.20	0.020923	45	66	46	268	272	0.35	0.11	3.3	3.92	9
310	1342229809	UGC11041	17:54:51.82	34:46:34.20	0.016332	31	19	341	239	187	0.58	0.08	3.3	3.08	10
311	1342231759	CGCG141-034	17:56:56.65	24:01:02.00	0.020162	51	65	450	384	370	0.39	0.18	3.3	3.78	9
312	1342239716	IRAS17578-0400	18:00:31.86	-04:00:53.40	0.013425	56	89	214	377	297	0.68	0.19	3.3	2.54	10
313	1342229821	IRAS18090+0130b	18:11:33.41	01:31:42.40	0.029190	-3	18	252	217	435	0.52	0.15	3.3	5.43	10
314	1342243548	IRAS18090+0130a	18:11:38.42	01:31:40.30	0.028936	-53	-23	408	97	217	0.61	0.12	3.3	5.37	10
315	1342225996	NGC6621	18:12:55.39	68:21:48.20	0.020843	84	63	299	318	348	0.56	0.08	3.3	3.91	10
316	1342230139	CGCG142-034b	18:16:33.83	22:06:38.90	0.018463	-44	44	448	442	453	0.48	0.27	3.3	3.47	10
317	1342230141	CGCG142-034a	18:16:40.69	22:06:46.40	...	...	...	...	455	517	0.51	0.13	3.5	3.51	10
318	1342222100	H1821+643	18:21:57.31	64:20:36.40	...	...	...	...	612	561	<0.002	2.96	3.3	41.24	9
319	1342192112	IRASF18293-3413	18:32:41.10	-34:11:27.00	0.018265	39	-22	287	383	391	0.63	0.24	3.4	3.43	6
320	1342233700	NGC6670Ab	18:33:34.26	59:53:17.90	0.028937	-34	-37	249	315	385	0.63	0.14	3.5	5.37	10
321	1342233698	NGC6670Aa	18:33:37.74	59:53:22.80	0.029141	43	26	121	262	260	0.61	0.27	3.3	5.42	10
322	1342192143	F18325-5926	18:36:58.29	-59:24:08.60	0.020101	27	62	450	599	767	0.02	1.18	3.3	3.77	1
323	1342251096	ESO103-G035	18:38:20.30	-65:25:40.00	0.013499	163 ^q	149	123	603	630	<0.003	1.44	3.4	2.55	14
324	1342241710	IC4734	18:38:25.75	-57:29:25.40	0.015496	-56	-34	201	341	453	0.38	0.1	3.3	2.92	9
325	1342232253	IRAS18443+7433	18:42:55.60	74:36:26.10	...	...	...	...	475	637	<0.051	0.21	2.3	22.19	9
326	1342235677	NGC6701	18:43:12.52	60:39:11.60	0.013181	-25	13	100	245	223	0.55	0.1	3.3	2.49	10
327	1342240169	ESO140-G043	18:44:53.98	-62:21:53.40	0.014195	-16	60	151	257	249	0.02	1.29	3.3	2.69	9
328	1342231316	1H1836-786	18:47:03.20	-78:31:51.00	0.074745	161 ^q	142	289	413	284	<0.012	1	3.3	13.17	9
329	1342231709	NGC6786b	19:10:54.01	73:24:36.00	0.025157	22	52	180	142	178	0.64	0.11	3.5	4.70	10
330	1342231707	NGC6786a	19:11:04.37	73:25:32.50	0.025319	12	23	198	225	292	0.29	0.24	3.3	4.72	10
331	1342231749	ESO593-IG008	19:14:31.15	-21:19:06.30	0.048834	-189 ^q	-328 ^r	496	667 ^o	352	0.56	0.19	3.5	8.87	9
332	1342231317	ESO-141-G055	19:21:14.15	-58:40:13.10	0.037265	17	91	100	279	330	<0.003	2.03	3.3	6.86	9
333	1342231722	IRAS19254-7245	19:31:21.55	-72:39:22.00	0.061594	-57	-151 ^r	576 ⁿ	835 ^o	1055 ^p	<0.032	0.37	2.3	11.01	8
334	1342208891	IRAS19297-0406	19:32:21.25	-03:59:56.30	...	...	...	...	460	349	0.3	0.13	4.2	14.92	1
335	1342208916	IRAS19542+1110	19:56:35.78	11:19:04.90	0.062666	82	89	273	335	629	0.29	0.27	2.3	11.20	1
336	1342232296	ESO339-G011	19:57:37.60	-37:56:08.40	0.019191	13	-19	350	306	331	0.27	0.94	3.3	3.61	9
337	1342232297	IRAS20037-1547	20:06:31.70	-15:39:08.00	0.192243	-45	8	138	244	200	0.04	0.2	3.3	29.75	9
338	1342232295	NGC6860	20:08:46.89	-61:06:00.70	0.014790	36	29	301	300	354	0.08	1.97	3.3	2.80	9
339	1342232302	IRAS20087-0308	20:11:23.86	-02:59:50.80	0.105416	-14	19	555 ⁿ	597	438	0.24	0.11	3.3	17.94	8
340	1342216371	IRAS20100-4156	20:13:29.85	-41:47:34.70	0.129707	-21	-11	216	433	478	<0.144	0.3	3.3	21.49	8
341	1342217702	NGC6907a	20:25:06.58	-24:48:32.90	0.010743	-10	20	270	172	100	0.57	0.06	3.3	2.04	10
342	1342232579	MCG+04-48-002a	20:28:35.07	25:44:00.30	0.014071	8	48	309	480	450	0.57	0.24	3.3	2.66	10
343	1342217654	NGC6926a	20:33:06.13	-02:01:38.90	0.019552	-141 ^q	-131 ^r	97	474	583	0.37	0.89	3.5	3.68	10
344	1342232581	IRAS20351+2521	20:37:17.73	25:31:37.50	0.034171	45	10	232	216	200	0.57	0.17	3.3	6.30	10
345	1342243654	Mkn509	20:44:09.74	-10:43:24.50	0.034585	46	55	399	563	364	0.02	1.22	3.3	6.39	14
346	1342217908	IRAS20414-1651	20:44:18.18	-16:40:16.40	0.087027	101	148	492	518	443	0.57	0.19	3.3	15.12	8
347	1342254215	IC5063-1	20:52:02.10	-57:04:06.60	0.011153	14	-37	396	422	395	<0.002	2.77	3.3	2.11	4
348	1342218619	CGCG448-020	20:57:24.38	17:07:39.20	0.036400	-19	-31	240	274	198	0.27	0.97	3.3	6.70	10
349	1342208934	IRAS20551-4250	20:58:26.78	-42:39:01.60	0.04296	117	88	177	226	111	0.06	0.2	3.3	7.86	1
350	1342232563	ESO286-G035	21:04:11.11	-43:35:36.10	0.017501	36	75	250	206	181	0.69	0.12	3.3	3.30	9
351	1342221628	IRAS21101+5810	21:11:29.28	58:23:07.90	0.039436	92	56	194	306	231	0.55	0.09	2.3	7.23	10
352	1342217815	ESO343-IG013	21:36:10.53	-38:32:42.80	0.019093	-6	25	270	240	134	0.61	0.13	2.2	3.59	10

TABLE 1 — *Continued*

No.	Herschel AOR	Name ^a	R.A. ^b	Dec. ^b	z^c [C II]	$dV[\text{Ne II}]^d$ km s^{-1}	$dV[\text{Ne III}]^e$ km s^{-1}	FWHM ^f [C II]	FWHM ^g [Ne II]	FWHM ^h [Ne III]	EW ⁱ μm	Neon ^j ratio	spax ^k	size ^l kpc	ref ⁿ
353	1342218185	NGC7130/IC5135	21:48:19.50	-34:57:06.00	0.016263	45	107	100	278	577	0.3	0.4	3.3	3.07	14
354	1342218489	NGC7172	22:02:01.90	-31:52:11.00	0.009265	227 ^q	217 ^r	199	527	351	0.03	0.71	3.4	(1.54)	14
355	1342256925	FSC22017+0319	22:04:19.17	03:33:50.20	0.061526	-28	-26	95	319	169	...	2.3	3.3	11.00	17
356	1342232569	NGC7213	22:09:16.25	-47:10:00.00	0.005986	47	43	252	549	517	<0.005	0.56	3.3	(1.00)	9
357	1342231750	ESO467-G027	22:14:39.97	-27:27:50.30	0.017552	-30	0	287	206	114	0.62	0.11	3.3	3.31	10
358	1342257681	3C445	22:23:49.57	-02:06:12.40	...	...	...	...	423	299	<0.002	1.54	3.3	10.06	17
359	1342233481	ESO602-G025	22:31:25.48	-19:02:04.10	0.025297	73	123	393	225	468	0.39	0.18	3.3	4.72	9
360	1342197892	NGC7314	22:35:46.20	-26:03:01.30	0.004739	20	36	100	126	192	<0.029	2.72	3.3	(0.83)	1
361	1342235675	UGC12138	22:40:17.05	08:03:14.10	0.025163	7	18	100	410	248	0.08	1.49	3.3	4.70	9
362	1342223714	UGC12150	22:41:12.26	34:14:57.00	0.021712	59	68	399	437	315	0.43	0.07	3.3	4.07	9
363	1342232568	ESO239-IG002	22:49:39.84	-48:50:58.30	0.043002	-15	20	295	396	410	0.31	0.1	3.3	7.86	9
364	1342211825	IRAS22491-1808	22:51:49.35	-17:52:24.00	0.077697	100	46	102	230	187	0.48	0.31	3.3	13.64	1
365	1342211171	NGC7469	23:03:15.62	08:52:26.40	0.016499	55	126	200	359	307	0.23	0.31	3.3	3.12	1
366	1342235676	Zw453.062	23:04:56.53	19:33:08.00	0.024931	-14	-8	248	271	384	0.58	0.4	3.3	4.66	9
367	1342235673	IRAS23060+0505	23:08:33.97	05:21:29.80	...	...	...	...	570	481	0.01	0.86	3.3	27.33	9
368	1342234062	Sersic159-03	23:13:58.80	-42:43:38.00	...	...	...	...	293	1234 ^p	...	1.48	3.3	10.42	13
369	1342210395	IRAS23128-5919	23:15:47.01	-59:03:16.90	0.044720	169 ^q	127	178	299	426	0.31	0.7	2.3	8.15	1
370	1342237590	IC5298	23:16:00.70	25:33:24.10	0.027363	15	139	252	375	542	<0.142	0.49	3.3	5.09	10
371	1342235751	NGC7591	23:18:16.28	06:35:08.90	0.016472	-6	-4	323	296	333	0.48	0.09	3.3	3.11	10
372	1342234259	NGC7592	23:18:22.60	-04:24:58.00	0.024721	29	67	200	392	392	0.3	0.24	2.3	4.62	10
373	1342211202	NGC7582	23:18:23.62	-42:22:13.60	0.005399	-59	-28	313	262	246	0.26	0.87	3.3	(0.97)	1
374	1342221357	NGC7582	23:18:23.62	-42:22:13.60	0.005467	-39	-8	307	262	246	0.26	0.87	3.3	(0.97)	14
375	1342222576	NGC7603	23:18:56.62	00:14:38.20	0.029413	-4	-26	398	402	498	0.02	0.49	3.3	5.47	9
376	1342232564	ESO077-IG014a	23:21:03.73	-69:13:01.00	0.041455	2	79	397	348	478	0.63	0.15	3.3	7.59	10
377	1342232566	ESO077-IG014b	23:21:05.45	-69:12:47.30	...	...	...	...	365	532	0.49	0.16	3.5	7.61	10
378	1342234997	IRAS23230-6926	23:26:03.57	-69:10:20.30	0.106342	-22	11	285	387	295	0.2	0.27	3.3	18.07	8
379	1342234264	NGC7674	23:27:56.72	08:46:44.50	0.028955	-15	8	197	400	559	0.02	1.95	3.3	5.37	10
380	1342235703	IRAS23253-5415	23:28:06.10	-53:58:31.00	0.129727	-101	-31	410	586	583	0.15	0.22	3.3	21.49	8
381	1342235842	NGC7679a	23:28:46.62	03:30:41.40	0.017193	-7	16	106	312	413	0.64	0.49	3.3	3.24	10
382	1342212520	Mkn930	23:31:58.30	28:56:50.00	0.018269	39	21	100	197	190	0.14	7.13	3.3	3.43	5
383	1342212515	IRAS23365+3604	23:39:01.29	36:21:09.80	0.064480	164 ^q	145	174	268	100	0.27	0.08	3.3	11.49	1
384	1342235844	MCG-01-60-022	23:42:00.91	-03:36:54.40	0.023165	-39	27	299	265	231	0.5	0.14	3.3	4.33	9
385	1342236288	IRAS23436+5257	23:46:05.43	53:14:01.60	0.034144	2	46	185	247	375	0.37	0.12	3.3	6.30	10
386	1342237595	NGC7752b	23:46:58.62	29:27:32.00	0.016642	26	-5	115	314	274	0.72	0.22	3.4	3.14	10
387	1342237593	NGC7752a	23:47:04.84	29:29:00.50	0.017321	21	46	235	294	336	0.32	0.19	4.2	3.26	10
388	1342236876	CGCG381-051	23:48:41.72	02:14:23.10	0.030934	1	58	196	338	602	0.54	0.13	3.3	5.73	9
389	1342238169	NGC7771c	23:51:04.00	20:09:02.00	0.014213	99	108	272	300	229	0.39	0.31	3.3	2.69	10
390	1342197840	Mkn331	23:51:26.80	20:35:10.00	0.018061	50	57	334	332	343	0.63	0.12	3.3	3.40	6

TABLE 1 — *Continued*

No.	Herschel AOR	Name ^a	R.A. ^b	Dec. ^b	z^c [C II]	$dV[\text{Ne II}]^d$ km s ⁻¹	$dV[\text{Ne III}]^e$ km s ⁻¹	FWHM ^f [C II]	FWHM ^g [Ne II]	FWHM ^h [Ne III]	EW ⁱ μm	Neon ^j ratio	spax ^k	size ^l kpc	ref ⁿ
-----	-----------------	-------------------	-------------------	-------------------	-----------------	--	---	-----------------------------	------------------------------	-------------------------------	-----------------------	----------------------------	-------------------	--------------------------	------------------

NOTE. —

^aSource name as used in CASSIS and the *Spitzer* Astronomical Observing Request.^bJ2000 coordinates of the source observed with the IRS short high aperture of size 4.7'' by 11.3''.^cRedshift z of [C II] emission line assuming rest wavelength of 157.74090 μm and corrected to Local Standard of Rest by HIPE.^dRadial velocity of [C II] emission line in km s⁻¹ relative to [Ne II] line, $dV = c[z([\text{C II}]) - z([\text{Ne II}])]$, assuming rest wavelength of 12.81355 μm for [Ne II] and corrected to Local Standard of Rest by adding VLSRCORR correction from headers in *Spitzer* AOR.^eRadial velocity of [C II] emission line in km s⁻¹ relative to [Ne III] line, $dV = c[z([\text{C II}]) - z([\text{Ne III}])]$, assuming rest wavelength of 15.55510 μm for [Ne III] and corrected to Local Standard of Rest by adding VLSRCORR correction from headers in *Spitzer* AOR.^fIntrinsic FWHM in km s⁻¹ of Gaussian fit to [C II] 158 μm emission line after quadratic correction for instrumental resolution having FWHM of 236 km s⁻¹. Gaussian fits to observed profiles having FWHM < instrumental resolution are arbitrarily assigned intrinsic FWHM of 100 km s⁻¹. Measured FWHM have typical uncertainties of ~ 1%; observed profiles with uncertainties in FWHM are shown by source name or position in cassis.sirtf.com/herschel.^gIntrinsic FWHM in km s⁻¹ of Gaussian fit to [Ne II] 12.81 μm emission line from CASSIS optimal extraction after quadratic correction for instrumental resolution having FWHM of 331 km s⁻¹. Gaussian fits to observed profiles having FWHM < instrumental resolution are arbitrarily assigned intrinsic FWHM of 100 km s⁻¹. Measured FWHM have typical uncertainties of ± 50 km s⁻¹, based on comparisons of independent spectral extractions. Observed profiles can be seen in cassis.sirtf.com.^hIntrinsic FWHM in km s⁻¹ of Gaussian fit to [Ne III] 15.55 μm emission line from CASSIS optimal extraction after quadratic correction for instrumental resolution having FWHM of 377 km s⁻¹. Gaussian fits to observed profiles having FWHM < instrumental resolution are arbitrarily assigned intrinsic FWHM of 100 km s⁻¹. Measured FWHM have typical uncertainties of ± 50 km s⁻¹, based on comparisons of independent spectral extractions. Observed profiles can be seen in cassis.sirtf.com.ⁱObserved equivalent width of 6.2 μm PAH feature from IRS low resolution spectra. EW defines source classification, with AGN having $\text{EW}(6.2 \mu\text{m}) < 0.1 \mu\text{m}$, composite (AGN plus starburst) having $0.1 \mu\text{m} < \text{EW}(6.2 \mu\text{m}) < 0.4 \mu\text{m}$, and starbursts having $\text{EW}(6.2 \mu\text{m}) > 0.4 \mu\text{m}$. EW measures are given by Sargsyan et al. (2011) or Stierwalt et al. (2013) or new measures from CASSIS if not in either reference.^jRatio of line fluxes for [Ne III]/[Ne II] for optimal extractions.^kPACS spaxel corresponding to position of IRS slit, using numbering convention in Figure 1. Overlays of PACS spaxels and IRS slits on source images are illustrated by source name or position in cassis.sirtf.com/herschel.^lProjected size of 9.4'' PACS spaxel in kpc at distance of source. Angular size distances determined from Wright (2006) for $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.27$ and $\Omega_\Lambda = 0.73$ (using [Ne II] redshift if [C II] redshift not listed) but adopting redshift independent distances (in parentheses) when given in the NASA/IPAC Extragalactic Database (NED).^mReference for *Herschel* PACS observing program that is source of data: 1 = SDP-esturm-3, KPGT-esturm-1 and DDT-esturm-4; 2 = OT2-idelooze-1; 3 = OT2-lsargsyan-1; 4 = OT2-pguillar-7; 5 = KPGT-smadde01-1; 6 = KPOT-pvanderw-1; 7 = OT1-bweiner-1; 8 = OT1-dfarrah-1 and OT2-dfarrah-5; 9 = OT1-dweedman-1; 10 = OT1-larmus-1; 11 = OT1-lyoung-1; 12 = OT1-nwerner-1; 13 = OT1-sveilleu-2; 14 = GT1-lspinogl-4 and GT2-lspinogl-6; 15 = KPOT-aedge-1; 16 = OT2-nnesvadb-3; 17 = OT2-tdiazsan-1.ⁿSource with intrinsic [C II] FWHM > 518 km s⁻¹, or more than 2 σ above median FWHM.^oSource with intrinsic [Ne II] FWHM > 631 km s⁻¹, or more than 2 σ above median FWHM.^pSource with intrinsic [Ne III] FWHM > 769 km s⁻¹, or more than 2 σ above median FWHM.^qSource with $dV([\text{C II}] - [\text{Ne II}]) > 131 \text{ km s}^{-1}$, or $dV([\text{C II}] - [\text{Ne II}]) < -63 \text{ km s}^{-1}$, values more than 2 σ above or below median difference.^rSource with $dV([\text{C II}] - [\text{Ne III}]) > 160 \text{ km s}^{-1}$, or $dV([\text{C II}] - [\text{Ne III}]) < -56 \text{ km s}^{-1}$, values more than 2 σ above or below median difference.